Index to Volume 88

Author Index

A

Abbott JH. Letter to the editor. [RE: Intertester reliability and validity of motion assessments during lumbar spine accessory motion testing.] 2008;88:536.

Able C: See Marchese VG

Addison O: See Marcus RL

Albers B: See Dunning K

Albert SF: See Boulton AJM

Alnwick GM. Misdiagnosis of serotonin syndrome as fibromyalgia and the role of physical therapists. 2008;88: 757–765.

Andersen CH: See Andersen LL

Andersen LL, Kjær M, Andersen CH, et al. Muscle activation during selected strength exercises in women with chronic neck muscle pain. 2008;88:703–711.

Anderson A: See Greenfield BH

Andrews AM: See Schnall BL

Anger MA: See Godges JJ

Angulo-Barroso RM: See Ulrich DA

Araújo LD: See Candotti CT

Ariail A, Sears T, Hampton E. Use of transabdominal ultrasound imaging in retraining the pelvic-floor muscles of a woman postpartum. 2008;88: 1208–1217.

Armijo Olivo S, Macedo LG, Gadotti IC, et al. Scales to assess the quality of randomized controlled trials: a systematic review. 2008;88:156–175.

Armstrong DG: See Boulton AJM

Asche CV: See Kirkness CS

Ashe MC: See Liu-Ambrose TY

B

Baccini M, Paci M, Nannetti L, et al. Scale for Contraversive Pushing: cutoff scores for diagnosing "pusher behavior" and construct validity. 2008;88:947–955.

Ballard DJ: See Dibble LE

Bandy WD: See Yates C

Banz R, Bolliger M, Colombo G, et al. Computerized visual feedback: an adjunct to robotic-assisted gait training. 2008;88:1135–1145.

Barbe MF: See Pierce SR

Barker KL: See Jenkins KL

Barr AE: See Pierce SR

Bart O: See Emanuel M

Bartlett DJ: See Hanna SE

Basso DM: See Herbert WJ

Baum BS: See Schnall BL

Beattie BL: See Liu-Ambrose TY

Beck LA, Einertson MJ, Winemiller MH, et al. Functional outcomes and quality of life after tumor-related hemipelvectomy. 2008;88:916–927.

Behrman AL, Nair PM, Bowden MG, et al. Author response to invited commentaries. [RE: Locomotor training restores walking in a nonambulatory child with chronic, severe, incomplete cervical spinal cord injury.] 2008;88:594–595.

Behrman AL, Nair PM, Bowden MG, et al. Locomotor training restores walking in a nonambulatory child with chronic, severe, incomplete cervical spinal cord injury. 2008;88: 580–590.

Behrman AL, Plummer-D'Amato P. Guest editorial: "What's in a name?" revisited. 2008;88:6–9.

Bell AL, Cavorsi J. Author response to invited commentary. [RE: Noncontact ultrasound therapy for adjunctive treatment of nonhealing wounds: retrospective analysis.] 2008;88:1526–1528.

Bell AL, Cavorsi J. Noncontact ultrasound therapy for adjunctive treatment of nonhealing wounds: retrospective analysis. 2008;88: 1517–1524.

Beneck GJ: See Powers CM

Bennett SE, Ohtake PJ. Letter to the editor. [RE: Journal publication productivity in academic physical therapy programs in the United States and Puerto Rico from 1998 to 2002.] 2008;88:539–540.

Bense B: See Miller EW

Berberich A: See Dunning K

Berg K: See Muir SW

Bergland A: See Ekerholt K

Biguet G: See Swärdh E

Billinger SA, Tseng BY, Kluding PM. Modified total-body recumbent stepper exercise test for assessing peak oxygen consumption in people with chronic stroke. 2008;88:1188–1195.

Binder-Macleod SA: See Chou LW

Birbaumer N: See Broetz D

Biricolti C: See Baccini M

Blair VP III: See Sinacore DR

Blasier RD: See Yates C

Blatt PJ: See Marchetti GF

Blum L, Korner-Bitensky N. Usefulness of the Berg Balance Scale in stroke rehabilitation: a systematic review. 2008;88:559–566.

Bohnert KL: See Hilton TN

Bohnert KL: See Mueller MJ

Bohnert KL: See Sinacore DR

Bolliger M: See Banz R

Booten AR: See Marchese VG

Borbely M: See Deutsch JE

Borello-France DF, Downey PA, Zyczynski HM, Rause CR. Continence and quality-of-life outcomes 6 months following an intensive pelvic-floor muscle exercise program for female stress urinary incontinence: a randomized trial comparing low- and high-frequency maintenance exercise. 2008;88:1545–1553.

Bosch PR: See Tompkins J

Boulton AJM, Armstrong DG, Albert SF, et al. Reprint—Comprehensive foot examination and risk assessment: a report of the Foot Care Interest Group of the American Diabetes Association, with endorsement by the American Association of Clinical Endocrinologists. [Reprinted from *Diabetes Care*. 2008;31:1679–1685.] 2008;88:1436–1443.

Bowden MG: See Behrman AL

Bowen P: See Marchese VG

Boyd LA, Vidoni ED, Siengsukon CF. Multidimensional motor sequence learning is impaired in older but not younger or middle-aged adults. 2008; 88:351–362.

Brach JS, Talkowski JB, Strotmeyer ES, Newman AB. Diabetes mellitus and gait dysfunction: possible explanatory factors. 2008;88:1365–1374.

Brach JS: See Huang WNW

Brach JS: See Talkowski JB

Broetz D, Soekadar SR, Birbaumer N. Letter to the editor. [RE: A four-week, task-specific neuroprosthesis program for a person with no active wrist or finger movement because of chronic stroke.] 2008;88:970.

Brosseau L, Wells GA, Tugwell P, et al. Ottawa Panel Evidence-Based Clinical Practice Guidelines for Aerobic Fitness Exercises in the Management of Fibromyalgia: Part 1. 2008;88:857–871.

Brosseau L, Wells GA, Tugwell P, et al. Ottawa Panel Evidence-Based Clinical Practice Guidelines for Strengthening Exercises in the Management of Fibromyalgia: Part 2. 2008;88:873–886.

Brownson RC: See Deshpande AD

Bulstra SK: See van den Akker-Scheek I

Bulstra SK: See Wagenmakers R

Burch A. Health care providers' knowledge, attitudes, and self-efficacy for working with patients with spinal cord injury who have diverse sexual orientations. 2008;88:191–198.

Burch A: See Miller EW

Burtner PA: See Sullivan KJ

Busch A: See Brosseau L

Busse ME, Khalil H, Quinn L, Rosser AE. Physical therapy intervention for people with Huntington disease. 2008;88:820–831.

C

Cade WT. Diabetes-related microvascular and macrovascular diseases in the physical therapy setting. 2008;88:1322–1335.

Campo MA. Author response to letter to the editor. [RE: Work-related musculoskeletal disorders in physical therapists: a prospective cohort study with 1-year follow-up.] 2008;88:887–888.

Campo M, Weiser S, Koenig KL, Nordin M. Work-related musculoskeletal disorders in physical therapists: a prospective cohort study with 1-year follow-up. 2008;88:608–619.

Candotti CT, Loss JF, Pressi AMS, et al. Electromyography for assessment of pain in low back muscles. 2008;88:1061–1067.

Casimiro L: See Brosseau L

Castro FAS: See Candotti CT

Cavorsi J: See Bell AL

Ceballos-Baumann A: See Starrost K

Cernak K, Stevens V, Price R, Shumway-Cook A. Locomotor training using body-weight support on a treadmill in conjunction with ongoing physical therapy in a child with severe cerebellar ataxia. 2008;88:88–97.

Cheeks JM: See Ross MD

Chen CN, Chuang LM, Wu YT. Clinical measures of physical fitness predict insulin resistance in people at risk for diabetes. 2008;88:1355–1364.

Chenowith R: See Tompkins J

Chesworth B: See Muir SW

Cheung RTH, Ng GYF. Influence of different footwear on force of landing during running. 2008;88:620–628.

Chou LW, Kesar TM, Binder-Macleod SA. Using customized rate-coding and recruitment strategies to maintain forces during repetitive activation of human muscles. 2008;88:363–375.

Christensen J: See Dibble LE

Christie A: See Jamtvedt G

Christie A: See Moe RH

Chuang LM: See Chen CN

Cieza A: See Geyh S

Cieza A: See Starrost K

Cohn JF: See Denlinger RL

Cohn JF: See Schmidt KL

Cohn R. Economic realities associated with diabetes care: opportunities to expand delivery of physical therapist services to a vulnerable population. 2008;88:1417–1424.

Colombo G: See Banz R

Combs SA: See Miller EW

Commean PK: See Hastings MK

Conn VS: See LeMaster JW

Connelly DM: See Mendelsohn ME

Connolly BH: See Marchese VG

Copeland JM, Taylor WJ, Dean SG. Factors influencing the use of outcome measures for patients with low back pain: a survey of New Zealand physical therapists. 2008;88:1492–1505.

Cosentino E: See Scivoletto G

Cox B: See Greenfield BH

Craik RL. Editorial: Climbing out of our silos to improve practice. 2008;88:555–558.

Craik RL. Editorial: Does clinical education need a *series* of tools to assess success? 2008;88:1106–1108.

Craik RL. Editorial: The 2008/2009 editorial board. 2008;88:986–988.

Craik RL, Harris S, Helders PJ. Editorial: In memoriam: Beverly Petterson Bishop (1922–2008). 2008;88:1454–1456.

Craik RL, Maher C. Editorial: Still "not satisfied". . . yet. 2008;88:423–425.

Craik RL: See Mangione KK

Crow JB, Gelfand B, Su EP. Use of joint mobilization in a patient with severely restricted hip motion following bilateral hip resurfacing arthroplasty. 2008;88:1591–1600.

Crow JL, Harmeling-van der Wel BC. Hierarchical properties of the motor function sections of the Fugl-Meyer Assessment Scale for people after stroke: a retrospective study. 2008;88:1554–1567.

D

Dahm KT: See Jamtvedt G

Dahm KT: See Moe RH

Dauser RC: See Behrman AL

Dean SG: See Copeland JM

Delgado-Charro MB: See Sylvestre JP

Delitto A. Thirty-Ninth Mary McMillan Lecture: We are what we do. 2008;88:1219–1227.

Delitto A: See Godges JJ

Denlinger RL, VanSwearingen JM, Cohn JF, Schmidt KL. Puckering and blowing facial expression in people with facial movement disorders. 2008;88:909-915.

Denlinger RL: See Schmidt KL

DePiero TJ: See Ellis T

DePompolo RW: See Beck LA

Deshpande AD, Dodson EA, Gorman I, Brownson RC. Physical activity and diabetes: opportunities for prevention through policy. 2008;88:1425-1435.

Deshpande AD, Harris-Hayes M, Schootman M. Epidemiology of diabetes and diabetes-related complications. 2008;88:1254-1264.

Deutsch JE, Borbely M, Filler J, et al. Use of a low-cost, commercially available gaming console (Wii) for rehabilitation of an adolescent with cerebral palsy. 2008;88:1196-1207.

Deutscher D, Hart DL, Dickstein R, et al. Implementing an integrated electronic outcomes and electronic health record process to create a foundation for clinical practice improvement. 2008;88:270-285.

Dibble LE, Christensen J, Ballard DJ, Foreman KB. Diagnosis of fall risk in Parkinson disease: an analysis of individual and collective clinical balance test interpretation. 2008; 88:323-332.

Dibble LE, Foreman KB, Ballard DJ, Christensen J. Author response to letter to the editor. [RE: Diagnosis of fall risk in Parkinson disease: an analysis of individual and collective clinical balance test interpretation.] 2008;88:679-680.

Dibble LE: See Marcus RL

Dickstein R: See Deutscher D

Dietz V: See Banz R

Di Fabio RP, Zampieri C, Tuite P. Gaze control and foot kinematics during stair climbing: characteristics leading to fall risk in progressive supranuclear palsy. 2008;88: 240-250.

Di Fabio RP: See Zampieri C

Dodd CAF: See Jenkins C

Dodson EA: See Deshpande AD

Downey PA: See Borello-France DF

Dubouloz CJ: See Brosseau L

Dudley-Javoroski S, Shields RK. Dose estimation and surveillance of mechanical loading interventions for bone loss after spinal cord injury. 2008;88:387-396.

Dunning K, Berberich A, Albers B, et al. A four-week, task-specific neuroprosthesis program for a person with no active wrist or finger movement because of chronic stroke. 2008;88:397-405.

Dunning K, Hill Hermann V, Levine PG. Author response to letter to the editor. [RE: A four-week, task-specific neuroprosthesis program for a person with no active wrist or finger movement because of chronic stroke.] 2008;88:970-972.

E

Edgerton VR. Invited commentary. [RE: Locomotor training restores walking in a nonambulatory child with chronic, severe, incomplete cervical spinal cord injury.] 2008;88: 590-591.

Editorial Office. Editorial: New author service: PTJ makes PubMed Central deposits. 2008;88:699-700.

Egan M: See Brosseau L

Einertson MJ: See Beck LA

Ekerholt K, Bergland A. Breathing: a sign of life and a unique area for reflection and action. 2008;88: 832-840.

Elkins MR: See Maher CG

Elboim M: See Laufer Y

Ellis T, Katz DI, White DK, et al. Effectiveness of an inpatient multidisciplinary rehabilitation program for people with Parkinson disease. 2008;88:812-819.

Emanuel M, Jarus T, Bart O. Effect of focus of attention and age on motor acquisition, retention, and transfer: a randomized trial. 2008;88:251-260.

F

Farquhar SJ, Reisman DS, Snyder-Mackler L. Persistence of altered movement patterns during a sit-to-stand task 1 year following unilateral total knee arthroplasty. 2008;88: 567-579.

Fetters L, Scholz JP. Invited commentary. [RE: Time series analysis of spontaneous upper-extremity movements of premature infants with brain injuries.] 2008;88: 1034-1036.

Fielder FA: See Sinacore DR

Fiers MJ: See Smits-Engelsman BCM

Filler J: See Deutsch JE

Finestone HM: See Brosseau L

Fish C: See Miller EW

Fisher JS: See Turcotte LP

Fitzgerald GK: See Schmitt LC

Foreman KB: See Dibble LE

Foster NE. Author response to invited commentary. [RE: Physical therapists' use of therapeutic exercise for patients with clinical knee osteoarthritis in the United Kingdom: in line with current recommendations?] 2008;88:1123.

Foster NE: See Holden MA

Freburger JK, Mielenz TJ. Invited commentary. [RE: Predictors of physical therapy clinic performance in the treatment of patients with low back pain syndromes.] 2008;88: 1004-1006.

Fredericson M: See Landel R

Fredericson M: See Powers CM

Fritz JM. Invited commentary. [RE: Predictors of physical therapy clinic performance in the treatment of patients with low back pain syndromes.] 2008;88:1007-1009.

Fritz JM: See Kirkness CS

Frykberg RG: See Boulton AJM

Fuentes J: See Armijo Olivo S

G

Gadotti IC: See Armijo Olivo S

Galloway JC: See Heathcock JC

Gelber J: See Hastings MK

Gelfand B: See Crow JB

Geyh S, Stucki G, Cieza A. Author response to invited commentary. [RE: Interrater reliability of the Extended ICF Core Set for stroke applied by physical therapists.] 2008;88:854-856.

Geyh S: See Starrost K

Godges JJ, Anger MA, Zimmerman G, Delitto A. Effects of education on return-to-work status for people with fear-avoidance beliefs and acute low back pain. 2008;88:231–239.

Gollish JD: See Kennedy DM

Gorman I: See Deshpande AD

Graf P: See Liu-Ambrose TY

Greenfield BH, Anderson A, Cox B, Tanner MC. Meaning of caring to 7 novice physical therapists during their first year of clinical practice. 2008;88:1154–1166.

Grindley EJ, Zizzi SJ, Nasypany AM. Author response to invited commentary. [RE: Use of protection motivation theory, affect, and barriers to understand and predict adherence to outpatient rehabilitation.] 2008;88:1543–1544.

Grindley EJ, Zizzi SJ, Nasypany AM. Use of protection motivation theory, affect, and barriers to understand and predict adherence to outpatient rehabilitation. 2008;88:1529–1540.

Groothoff JW: See van den Akker-Scheek I

Groothoff JW: See Wagenmakers R

Groothuis-Oudshoorn CG: See Kottink AIR

Grunow J: See Starrost K

Guarrera-Bowlby P: See Deutsch JE

Gulve EA. Exercise and glycemic control in diabetes: benefits, challenges, and adjustments to pharmacotherapy. 2008;88:1297–1321.

Gutvirtz M: See Deutscher D

Guy RH: See Sylvestre JP

H

Haavardsholm E: See Jamtvedt G

Haavardsholm EA: See Moe RH

Hagen KB: See Jamtvedt G

Hagen KB: See Moe RH

Haines-Wangda A: See Brosseau L

Halili A. Letter to the editor. [RE: Puckering and blowing facial expression in people with facial movement disorders.] 2008;88:1231.

Hall D: See Schenkman M

Halloran SW. Letter to the editor. [RE: Work-related musculoskeletal disorders in physical therapists: a prospective cohort study with 1-year follow-up.] 2008;88:887.

Hampton E: See Ariail A

Hancock ML: See Marchese VG

Hanna SE, Bartlett DJ, Rivard LM, Russell DJ. Reference curves for the Gross Motor Function Measure: percentiles for clinical description and tracking over time among children with cerebral palsy. 2008;88:596–607.

Hanna SE: See Kennedy DM

Hansen K: See Andersen LL

Hansen PB: See Andersen LL

Harmeling-van der Wel BC: See Crow JL

Harris S: See Craik RL

Harris SR. Letter to the editor. [RE: Journal publication productivity in academic physical therapy programs in the United States and Puerto Rico from 1998 to 2002.] 2008;88:791.

Harris SR: See Keays KS

Harris-Hayes M: See Deshpande AD

Hart DL: See Deutscher D

Hart DL: See Resnik L

Harth M: See Brosseau L

Hastings MK, Gelber J, Commean PK, et al. Bone mineral density of the tarsals and metatarsals with reloading. 2008;88:766–779.

Hastings MK: See Sinacore DR

Hay EM: See Holden MA

Hay EM: See Wood L

Heathcock JC, Lobo M, Galloway JC. Movement training advances the emergence of reaching in infants born at less than 33 weeks of gestational age: a randomized clinical trial. 2008;88:310–322.

Heiss DG: See Herbert WJ

Helders PJ: See Craik RL

Helgeson K, Smith AR Jr. Process for applying the *International Classification of Functioning, Disability and Health* model to a patient with patellar dislocation. 2008;88:956–964.

Hellesnes B: See Nilsson BB

Hellman R: See Boulton AJM

Henderson L: See Smits-Engelsman BCM

Henderson SE: See Smits-Engelsman BCM

Herbert RD: See Maher CG

Herbert WJ, Heiss DG, Basso DM. Influence of feedback schedule in motor performance and learning of a lumbar multifidus muscle task using rehabilitative ultrasound imaging: a randomized clinical trial. 2008;88:261–269.

Herget BR: See Behrman AL

Hermens HJ: See Kottink AIR

Hetland ML: See Holm B

Hill Hermann VA: See Dunning K

Hilton TN, Tuttle LJ, Bohnert KL, et al. Excessive adipose tissue infiltration in skeletal muscle in individuals with obesity, diabetes mellitus, and peripheral neuropathy: association with performance and function. 2008;88:1336–1344.

Hohler AD: See Ellis T

Holden MA, Nicholls EE, Hay EM, Foster NE. Physical therapists' use of therapeutic exercise for patients with clinical knee osteoarthritis in the United Kingdom: in line with current recommendations? 2008;88:1109–1121.

Holm B, Jacobsen S, Skjodt H, et al. Keitel Functional Test for patients with rheumatoid arthritis: translation, reliability, validity, and responsiveness. 2008;88:664–678.

Holm I: See Jamtvedt G

Hoppe KM: See Beck LA

Horn SD: See Deutscher D

Howard S: See Marchese VG

Howard SM: See Straub SJ

Howland DR: See Behrman AL

Huang WNW, VanSwearingen JM, Brach JS. Gait variability in older adults: observational rating validated by comparison with a computerized walkway gold standard. 2008;88:1146–1153.

Huhn K: See Deutsch JE

I

IJzerman MJ: See Kottink AIR

J

Jacobs ML: See Wagenmakers R

Jacobsen S: See Holm B

Jamtvedt G, Dahm KT, Christie A, et al. Physical therapy interventions for patients with osteoarthritis of the knee: an overview of systematic reviews. 2008;88:123–136.

Jamtvedt G: See Moe RH

Jan MH, Lin JJ, Liau JJ, et al. Investigation of clinical effects of high- and low-resistance training for patients with knee osteoarthritis: a randomized controlled trial. 2008;88: 427–436.

Jarus T: See Emanuel M

Jenkins C, Barker KL, Pandit H, et al. After partial knee replacement, patients can kneel, but they need to be taught to do so: a single-blind randomized controlled trial. 2008;88: 1012–1021.

Jensen T: See Holm B

Jette AM. Invited commentary. [RE: Interrater reliability of the Extended ICF Core Set for stroke applied by physical therapists.] 2008;88: 851–853.

Johns LD: See Straub SJ

Johnson JE: See Sinacore DR

Jones A: See Silva LE

K

Kantak SS: See Sullivan KJ

Kaste SC: See Marchese VG

Katz DI: See Ellis T

Keays KS, Harris SR, Lucyshyn JM, MacIntyre DL. Effects of Pilates exercises on shoulder range of motion, pain, mood, and upper-extremity function in women living with breast cancer: a pilot study. 2008;88:494–510.

Kennedy DM, Stratford PW, Riddle DL, et al. Assessing recovery and establishing prognosis following total knee arthroplasty. 2008;88:22–32.

Kesar TM: See Chou LW

Khalil H: See Busse ME

Khan KM: See Liu-Ambrose TY

King WM: See Pandya S

Kirkman MS: See Boulton AJM

Kirkness CS, Marcus RL, LaStayo PC, et al. Diabetes and associated risk factors in patients referred for physical therapy in a national primary care electronic medical record database. 2008;88: 1408–1416.

Kissela BM: See Page SJ

Kjær M: See Andersen LL

Klarlund M: See Holm B

Kluding PM: See Billinger SA

Koenig KL: See Campo M

Kohrt WM: See Schenkman M

Korner-Bitensky N: See Blum L

Kottink AIR, Hermens HJ, Nene AV, et al. Therapeutic effect of an implantable peroneal nerve stimulator in subjects with chronic stroke and footdrop: a randomized controlled trial. 2008;88:437–448.

Krieger MM: See Richter RR

Kulig K: See Landel R

Kulig K: See Powers CM

Kumar R: See Schenkman M

L

Lambert K: See Brosseau L

Landel R, Kulig K, Fredericson M, et al. Author response to letter to the editor. [RE: Intertester reliability and validity of motion assessments during lumbar spine accessory motion testing.] 2008;88:536–537.

Landel R, Kulig K, Fredericson M, et al. Intertester reliability and validity of motion assessments during lumbar spine accessory motion testing. 2008;88:43–49.

Landel RF: See Powers CM

LaStayo PC: See Kirkness CS

LaStayo PC: See Marcus RL

La Torre M: See Candotti CT

Laufer Y, Elboim M. Effect of burst frequency and duration of kilohertz-frequency alternating currents and of low-frequency pulsed currents on strength of contraction, muscle fatigue, and perceived discomfort. 2008;88:1167–1176.

Lauer RT: See Pierce SR

Lavery LA: See Boulton AJM

Leeb BF, Rintelen B. Letter to the editor. [RE: Effectiveness of nonpharmacological and nonsurgical interventions for hip osteoarthritis: an umbrella review of high-quality systematic reviews.] 2008;88: 407–408.

Leemrijse CJ, Swinkels ICS, Veenhof C. Direct access to physical therapy in the Netherlands: results from the first year in community-based physical therapy. 2008;88:936–946.

LeMaster JW, Mueller MJ, Reiber GE, et al. Effect of weight-bearing activity on foot ulcer incidence in people with diabetic peripheral neuropathy: Feet First randomized controlled trial. 2008;88:1385–1398.

LeMaster JW: See Boulton AJM

Leonard A: See Page SJ

Leskinen E: See Paltamaa J

Levine P: See Page S

Levine P: See Page SJ

Levine PG: See Dunning K

Li B: See Landel R

Liau JJ: See Jan MH

Lin DH: See Jan MH

Lin JJ: See Jan MH

Lin YF: See Jan MH

Liu D: See Resnik L

Liu-Ambrose TY, Ashe MC, Graf P, et al. Increased risk of falling in older community-dwelling women with mild cognitive impairment. 2008;88: 1482–1491.

Lloyd MC: See Ulrich DA

Lobo M: See Heathcock JC

Long TM, Perry DF. Pediatric physical therapists' perceptions of their training in assistive technology. 2008;88:629–639.

Loo KK: See Ohgi S

Looper JE: See Ulrich DA

Lopopolo RB: See Mangione KK

Loss JF: See Candotti CT

Lucyshyn JM: See Keays KS

Lund H. Invited commentary. [RE: Physical therapists' use of therapeutic exercise for patients with clinical knee osteoarthritis in the United Kingdom: in line with current recommendations?] 2008;88: 1121–1122.

Lünenburger L: See Banz R

M

Macedo LG: See Armijo Olivo S

MacIntyre DL: See Keays KS

MacIntyre DL: See Mathur S

Madsen RW: See LeMaster JW

Magee DJ: See Armijo Olivo S

Maher C. Letter to the editor. [RE: Journal publication productivity in academic physical therapy programs in the United States and Puerto Rico from 1998 to 2002.] 2008;88:539.

Maher C: See Craik RL

Maher CG, Moseley AM, Sherrington C, et al. A description of the trials, reviews, and practice guidelines indexed in the PEDro database. 2008;88:1068–1077.

Mälkiä E: See Paltamaa J

Mammone A: See Scivoletto G

Mangione KK, Lopopolo RB, Neff NP, et al. Interventions used by physical therapists in home care for people after hip fracture. 2008;88:199–210.

Marchese VG, Connolly BH, Able C, et al. Relationships among severity of osteonecrosis, pain, range of motion, and functional mobility in children, adolescents, and young adults with acute lymphoblastic leukemia. 2008;88:341–350.

Marchetti GF, Whitney SL, Blatt PJ, et al. Temporal and spatial characteristics of gait during performance of the Dynamic Gait Index in people with and people without balance or vestibular disorders. 2008;88:640–651.

Marcus RL, Smith S, Morrell G, et al. Comparison of combined aerobic and high-force eccentric resistance exercise with aerobic exercise only for people with type 2 diabetes mellitus. 2008;88:1345–1354.

Marcus RL: See Kirkness CS

Marshall AD: See Brosseau L

Martin JB: See Behrman AL

Mathur S, Takai KPR, MacIntyre DL, Reid D. Estimation of thigh muscle mass with magnetic resonance imaging in older adults and people with chronic obstructive pulmonary disease. 2008;88:219–230.

McEwen IR. Invited commentary. [RE: Locomotor training restores walking in a nonambulatory child with chronic, severe, incomplete cervical spinal cord injury.] 2008;88:592–594.

McGowan J: See Brosseau L

McMeeken JM: See Wilde VE

Mehr DR: See LeMaster JW

Melo MO: See Candotti CT

Mendelsohn ME, Connelly DM, Overend TJ, Petrella PJ. Validity of values for metabolic equivalents of task during submaximal all-extremity exercise and reliability of exercise responses in frail older adults. 2008;88:747–756.

Mielenz TJ: See Freburger JK

Miller EW, Combs SA, Fish C, et al. Running training after stroke: a single-subject report. 2008;88: 511–522.

Mills JL Sr: See Boulton AJM

Mizuike C: See Ohgi S

Moe RH, Haavardsholm EA, Christie A, et al. Author response to letter to the editor. [RE: Effectiveness of nonpharmacological and nonsurgical interventions for hip osteoarthritis: an umbrella review of high-quality systematic reviews.] 2008;88:408.

Moe RH: See Jamtvedt G

Moldofsky H: See Brosseau L

Molinari M: See Scivoletto G

Mor V: See Resnik L

Morita S: See Ohgi S

Morrell G: See Marcus RL

Morris LO: See Marchetti GF

Mortellite K: See Dunning K

Moseley AM: See Maher CG

Mossberg KA, Orlander EE, Norcross JL. Cardiorespiratory capacity after weight-supported treadmill training in patients with traumatic brain injury. 2008;88:77–87.

Mueller Mj. Editorial: People with diabetes: a population desperate for movement. 2008;88:1250–1253.

Mueller MJ, Zou D, Bohnert KL, et al. Plantar stresses on the neuropathic foot during barefoot walking. 2008;88:1375–1384.

Mueller MJ: See Boulton AJM

Mueller MJ: See Hilton TN

Mueller MJ: See LeMaster JW

Muir SW, Berg K, Chesworth B, Speechley M. Use of the Berg Balance Scale for predicting multiple falls in community-dwelling elderly people: a prospective study. 2008;88:449–459.

Murphy AJ: See Rees SS

Murray DW: See Jenkins C

Myamoto S: See Silva LE

N

Nair PM: See Behrman AL

Nannetti L: See Baccini M

Nasypany AM: See Grindley EJ

Natour J: See Silva LE

Neel MD: See Marchese VG

Neff NP: See Mangione KK

Nene AV: See Kottink AIR

Newman AB: See Brach JS

Newman AB: See Talkowski JB

Ng GYF: See Cheung RTH

Nicholls EE: See Holden MA

Nielson W: See Brosseau L

Nilsson BB, Hellesnes B, Westheim A, Risberg MA. Group-based aerobic interval training in patients with chronic heart failure: Norwegian Ullevaal Model. 2008;88:523–535.

Nitsch W: See Vaughn HT

Norcross JL: See Mossberg KA

Nordin M: See Campo M

Norton BJ. Letter to the editor. [RE: Diagnosis of fall risk in Parkinson disease: an analysis of individual and collective clinical balance test interpretation.] 2008;88:679.

O

Ohgi S, Morita S, Loo KK, Mizuike C. Author response to invited commentary. [RE: Time series analysis of spontaneous upper-extremity movements of premature infants with brain injuries.] 2008;88:1037–1038.

Ohgi S, Morita S, Loo KK, Mizuike C. Time series analysis of spontaneous upper-extremity movements of premature infants with brain injuries. 2008;88:1022–1033.

Ohtake PJ: See Bennett SE

Oliveira LM: See Silva LE

Opava CH: See Swärdh E

Orlander EE: See Mossberg KA

Ostergaard M: See Holm B

Overend TJ: See Mendelsohn ME

Owens A: See Miller EW

P

Paci M: See Baccini M

Page S, Levine P. Author response to letter to the editor. [RE: Modified constraint-induced therapy in chronic stroke: results of a single-blinded randomized controlled trial.] 2008;88:684–688.

Page SJ, Levine P, Leonard A, et al. Modified constraint-induced therapy in chronic stroke: results of a single-blinded randomized controlled trial. 2008;88:333–340.

Page SJ: See Dunning K

Palombaro KM: See Mangione KK

Paltamaa J, Sarasoja T, Leskinen E, et al. Measuring deterioration in International Classification of Functioning domains of people with multiple sclerosis who are ambulatory. 2008;88:176–190.

Pandit H: See Jenkins C

Pandya S, King WM, Tawil R. Facioscapulohumeral dystrophy. 2008;88:105–113.

Pasini M: See Candotti CT

Patten C: See Wagner JM

Peat G: See Wood L

Perry DF: See Long TM

Pessanha APC: See Silva LE

Petrella PJ: See Mendelsohn ME

Phadke CP: See Behrman AL

Pierce SR, Barbe MF, Barr AE, et al. Roles of reflex activity and co-contraction during assessments of spasticity of the knee flexor and knee extensor muscles in children with cerebral palsy and different functional levels. 2008;88:1124–1134.

Plummer-D'Amato P: See Behrman AL

Poitras S: See Brosseau L

Porter BM: See Marchese VG

Powers CM, Beneck GJ, Kulig K, et al. Effects of a single session of posterior-to-anterior spinal mobilization and press-up exercise on pain response and lumbar spine extension in people with nonspecific low back pain. 2008;88:485–493.

Powers CM: See Landel R

Pressi AMS: See Candotti CT

Price R: See Cernak K

Prior F: See Hastings MK

Prosiegel M: See Starrost K

Pui CH: See Marchese VG

Q

Quinn L: See Busse ME

R

Rai SN: See Marchese VG

Rause CR: See Borello-France DF

Rees SS, Murphy AJ, Watsford ML. Effects of whole-body vibration exercise on lower-extremity muscle strength and power in an older population: a randomized clinical trial. 2008;88:462–470.

Reiber GE: See LeMaster JW

Reid D: See Mathur S

Reier PJ: See Behrman AL

Reisman AS: See Schmitt LC

Reisman DS: See Farquhar SJ

Resnik L, Hart DL. Author response to invited commentaries. [RE: Predictors of physical therapy clinic performance in the treatment of patients with low back pain syndromes.] 2008;88:1009–1011.

Resnik L, Liu D, Hart DL, Mor V. Benchmarking physical therapy clinic performance: statistical methods to enhance internal validity when using observational data. 2008;88:1078–1087.

Resnik L, Liu D, Mor V, Hart DL. Predictors of physical therapy clinic performance in the treatment of patients with low back pain syndromes. 2008;88:989-1004.

Rhodes JA: See Wagner JM

Richter RR, Schlomer SL, Krieger MM, Siler WL. Author response to letters to the editor. [RE: Journal publication productivity in academic physical therapy programs in the United States and Puerto Rico from 1998 to 2002.] 2008;88:540–541.

Richter RR, Schlomer SL, Krieger MM, Siler WL. Journal publication productivity in academic physical therapy programs in the United States and Puerto Rico from 1998 to 2002. 2008;88:376–386.

Riddle DL: See Kennedy DM

Rinaldi LA: See Baccini M

Rintelen B: See Leeb BF

Risberg MA: See Nilsson BB

Rivard LM: See Hanna SE

Robertson VJ. Invited commentary. [RE: Noncontact ultrasound therapy for adjunctive treatment of nonhealing wounds: retrospective analysis.] 2008;88:1524–1526.

Robertson VJ. Invited commentary. [RE: Variability in effective radiating area at 1 MHz affects ultrasound treatment intensity.] 2008;88:58–61.

Robinson VA: See Brosseau L

Ross MD, Cheeks JM. Undetected hangman's fracture in a patient referred for physical therapy for the treatment of neck pain following trauma. 2008;88:98–104.

Rosser AE: See Busse ME

Rudolph KS: See Schmitt LC

Russell DJ: See Hanna SE

Russell-Doreleyers M: See Brosseau L

S

Saint-Hilaire M: See Ellis T

Sarasoja T: See Paltamaa J

Schenkman M, Hall D, Kumar R, Kohrt WM. Endurance exercise training to improve economy of movement of people with Parkinson disease: three case reports. 2008;88: 63–76.

Schlomer SL: See Richter RR

Schmidt KL, Denlinger RL, VanSwearingen JM, Cohn JF. Author response to letter to the editor. [RE: Puckering and blowing facial expression in people with facial movement disorders.] 2008;88: 1231–1232.

Schmidt KL: See Denlinger RL

Schmitt LC, Fitzgerald GK, Reisman AS, Rudolph KS. Instability, laxity, and physical function in patients with medial knee osteoarthritis. 2008;88:1506–1516.

Schnall BL, Baum BS, Andrews AM. Gait characteristics of a soldier with a traumatic hip disarticulation. 2008;88:1568–1577.

Scholz JP: See Fetters L

Schootman M: See Deshpande AD

Scivoletto G, Cosentino E, Mammone A, Molinari M. Inflammatory myelopathies and traumatic spinal cord lesions: comparison of functional and neurological outcomes. 2008;88:471–484.

Sears T: See Ariail A

Senesac CR: See Behrman AL

Seney M: See Steffen T

Sheehan P: See Boulton AJM

Sherrington C: See Maher CG

Shewokis PA: See Pierce SR

Shields RK: See Dudley-Javoroski S

Shumway-Cook A: See Cernak K

Siengsukon CF: See Boyd LA

Siler WL: See Richter RR

Silva LE, Valim V, Pessanha APC, et al. Hydrotherapy versus conventional land-based exercise for the management of patients with osteoarthritis of the knee: a randomized clinical trial. 2008;88: 12–21.

Sim FF: See Beck LA

Sim J: See Wood L

Sinacore DR, Hastings MK, Bohnert KL, et al. Inflammatory osteolysis in diabetic neuropathic (Charcot) arthropathies of the foot. 2008;88: 1399–1407.

Sinacore DR: See Hastings MK

Sinacore DR: See Hilton TN

Sinacore DR: See Mueller MJ

Sjøgaard G: See Andersen LL

Skjodt H: See Holm B

Smith AR Jr: See Helgeson K

Smith S: See Marcus RL

Smits-Engelsman BCM, Fiers MJ, Henderson SE, Henderson L. Interrater reliability of the Movement Assessment Battery for Children. 2008;88:286–294.

Snyder-Mackler L: See Farquhar SJ

Soekadar SR: See Broetz D

Speechley M: See Muir SW

Stanton T: See Armijo Olivo S

Starrost K, Geyh S, Trautwein A, et al. Interrater reliability of the Extended ICF Core Set for stroke applied by physical therapists. 2008;88: 841–851.

Steffen T, Seney M. Author response to letter to the editor. [RE: Test-retest reliability and minimal detectable change on balance and ambulation tests, the 36-Item Short-Form Health Survey, and the Unified Parkinson Disease Rating Scale in people with parkinsonism.] 2008;88:891.

Steffen T, Seney M. Test-retest reliability and minimal detectable change on balance and ambulation tests, the 36-Item Short-Form Health Survey, and the Unified Parkinson Disease Rating Scale in people with parkinsonism. 2008;88:733–746.

Stehno-Bittel L. Intricacies of fat. 2008;88:1265–1278.

Stevens M: See van den Akker-Scheek I

Stevens M: See Wagenmakers R

Stevens V: See Cernak K

Stratford PW. Letter to the editor. [RE: Test-retest reliability and minimal detectable change on balance and ambulation tests, the 36-Item Short-Form Health Survey, and the Unified Parkinson Disease Rating Scale in people with parkinsonism.] 2008;88:888–890.

Stratford PW: See Kennedy DM

Straub SJ, Johns LD, Howard SM. Author response to invited commentary. [RE: Variability in effective radiating area at 1 MHz affects ultrasound treatment intensity.] 2008;88:61–62.

Straub SJ, Johns LD, Howard SM. Variability in effective radiating area at 1 MHz affects ultrasound treatment intensity. 2008;88:50–57.

Strobl RM. Letter to the editor. [RE: Indicators of lumbar zygapophyseal joint pain: survey of an expert panel with the Delphi technique.] 2008;88:137.

Strotmeyer ES: See Brach JS

Stucki G: See Geyh S

Stucki G: See Starrost K

Studenski S: See Talkowski JB

Studenski SA. Invited commentary. [RE: Use of the Berg Balance Scale for predicting multiple falls in community-dwelling elderly people: a prospective study.] 2008;88: 460–461.

Su EP: See Crow JB

Sullivan KJ, Kantak SS, Burtner PA. Motor learning in children: feedback effects on skill acquisition. 2008;88: 720–732.

Swain JM: See Tompkins J

Swärdh E, Biguet G, Opava CH. Views on exercise maintenance: variations among patients with rheumatoid arthritis. 2008;88:1049–1060.

Swinkels ICS: See Leemrijse CJ

Sylvestre JP, Guy RH, Delgado-Charro MB. *In vitro* optimization of dexamethasone phosphate delivery by iontophoresis. 2008;88: 1177–1185.

Szaflarski JP: See Page SJ

T

Takai KPR: See Mathur S

Talkowski JB, Brach JS, Studenski S, Newman AB. Impact of health

perception, balance perception, fall history, balance performance, and gait speed on walking activity in older adults. 2008;88:1474–1481.

Talkowski JB: See Brach JS

Tanner MC: See Greenfield BH

Tawil R: See Pandya S

Taylor MJ. Letter to the editor. [RE: Breathing: a sign of life and a unique area for reflection and action.] 2008;88:972.

Taylor WJ: See Copeland JM

Tenniglo MJ: See Kottink AIR

Thomas E: See Wood L

Thompson FJ: See Behrman AL

Thorstensson CA. Invited commentary. [RE: Use of protection motivation theory, affect, and barriers to understand and predict adherence to outpatient rehabilitation.] 2008;88:1541–1543.

Tiede JL: See Tompkins J

Tiernan CW: See Ulrich DA

Tompkins J, Bosch PR, Chenowith R, et al. Changes in functional walking distance and health-related quality of life after gastric bypass surgery. 2008;88:928–935.

Trautwein A: See Starrost K

Tseng BY: See Billinger SA

Tugwell P: See Brosseau L

Tuite P: See Di Fabio RP

Turcotte LP, Fisher JS. Skeletal muscle insulin resistance: roles of fatty acid metabolism and exercise. 2008;88:1279–1296.

Tuttle LJ: See Hilton TN

Tuttle LJ: See Mueller MJ

U

Ulrich DA, Lloyd MC, Tiernan CW, et al. Effects of intensity of treadmill training on developmental outcomes and stepping in infants with Down syndrome: a randomized trial. 2008;88:114–122.

Underwood FB. Invited commentary. [RE: *In vitro* optimization of dexamethasone phosphate delivery by iontophoresis.] 2008;88:1185–1187.

V

Valim V: See Silva LE

Vance JM: See Marchetti GF

van den Akker-Scheek I, Zijlstra W, Groothoff JW, et al. Physical functioning before and after total hip arthroplasty: perception and performance. 2008;88:712–719.

van den Akker-Scheek I: See Wagenmakers R

VanSwearingen JM: See Denlinger RL

VanSwearingen JM: See Huang WNW

VanSwearingen JM: See Schmidt KL

Vaughn HT, Nitsch W. Ilial anterior rotation hypermobility in a female collegiate tennis player. 2008;88:1578–1590.

Veenhof C: See Leemrijse CJ

Veilleux L: See Brosseau L

Vidoni ED: See Boyd LA

Villareal DT: See Sinacore DR

W

Wagenmakers R, Stevens M, van den Akker-Scheek I, et al. Predictive value of the Western Ontario and McMaster Universities Osteoarthritis Index for the amount of physical activity after total hip arthroplasty. 2008;88:211–218.

Wagenmakers R, Stevens M, Zijlstra W, et al. Habitual physical activity behavior of patients after primary total hip arthroplasty. 2008;88:1039–1048.

Wagner JM, Rhodes JA, Patten C. Reproducibility and minimal detectable change of three-dimensional kinematic analysis of reaching tasks in people with hemiparesis after stroke. 2008;88:652–663.

Wahoff-Stice D: See Marcus RL

Ward RS. 2008 Presidential Address: Our great opportunity. 2008;88:1228–1230.

Warden SJ. Letter to the editor. [RE: Journal publication productivity in academic physical therapy programs in the United States and Puerto Rico from 1998 to 2002.] 2008;88:538–539.

Watsford ML: See Rees SS

Weiser S: See Campo M

Wells GA: See Brosseau L

Westheim A: See Nilsson BB

White DK: See Ellis T

Whitney SL: See Marchetti GF

Wikström J: See Paltamaa J

Wilde VE, McMeeken JM. Author response to letter to the editor. [RE: Indicators of lumbar zygapophyseal joint pain: survey of an expert panel with the Delphi technique.] 2008;88:137.

Wilson KG: See Brosseau L

Winemiller MH: See Beck LA

Wolf SL. Letter to the editor. [RE: Modified constraint-induced therapy in chronic stroke: results of a single-blinded randomized controlled trial.] 2008;88:680–684.

Wood L, Peat G, Thomas E, et al. Associations between physical examination and self-reported physical function in older community-dwelling adults with knee pain. 2008;88:33–42.

Wukich DK: See Boulton AJM

Wu YT: See Chen CN

Y

Yates C, Bandy WD, Blasier RD. Traumatic dislocation of the hip in a high school football player. 2008;88:780–788.

Z

Zampieri C, Di Fabio RP. Balance and eye movement training to improve gait in people with progressive supranuclear palsy: a quasi-randomized clinical trial. 2008;88:1460–1473.

Zampieri C: See Di Fabio RP

Zebis MK: See Andersen LL

Zijlstra W: See van den Akker-Scheek I

Zijlstra W: See Wagenmakers R

Zimmerman G: See Godges JJ

Zizzi SJ: See Grindley EJ

Zou D: See Mueller MJ

Zyczynski HM: See Borello-France DF

Subject Index

A

Administration

Records

Implementing an integrated electronic outcomes and electronic health record process to create a foundation for clinical practice improvement. Deutscher D, Hart DL, Dickstein R, et al. 2008;88:270-285.

Ambulation Aids

General

Influence of different footwear on force of landing during running. Cheung RTH, Ng GYF. 2008;88:620-628.

- Correction. 2008;88:797.

American Physical Therapy Association

Ethics and Judicial Committee Disciplinary Action. 2008;88:893.

Membership statistics. 2008;88:798.

Thirty-Ninth Mary McMillan Lecture: We are what we do. Delitto A. 2008;88:1219-1227.

2008 Presidential Address: Our great opportunity. Ward RS. 2008;88:1228-1230.

Arthritis

Osteoarthritis

Assessing recovery and establishing prognosis following total knee arthroplasty. Kennedy DM, Stratford PW, Riddle DL, et al. 2008;88:22-32.

[Effectiveness of nonpharmacological and nonsurgical interventions for hip osteoarthritis: an umbrella review of high-quality systematic reviews. Christie A, Jamtvedt G, Dahm KT, et al. 2007;87:1716-1727.]

- Letter to the editor. Leeb BF, Rintelen B. 2008;88:407-408.
- Author response. Moe RH, Haavardsholm EA, Christie A, et al. 2008;88:408.

Hydrotherapy versus conventional land-based exercise for the management of patients with osteoarthritis of the knee: a randomized clinical trial. Silva LE, Valim V, Pessanha APC, et al. 2008;88:12-21.

Instability, laxity, and physical function in patients with medial knee osteoarthritis. Schmitt LC, Fitzgerald GK, Reisman AS, Rudolph KS. 2008;88:1506-1516.

Investigation of clinical effects of high- and low-resistance training for patients with knee osteoarthritis: a randomized controlled trial. Jan MH, Lin JJ, Liau JJ, et al. 2008;88:427-436.

Physical therapists' use of therapeutic exercise for patients with clinical knee osteoarthritis in the United Kingdom: in line with current recommendations? Holden MA, Nicholls EE, Hay EM, Foster NE. 2008;88:1109-1121.

- Invited commentary. Lund H. 2008;88:1121-1122.
- Author response. Foster NE. 2008;88:1123.

Physical therapy interventions for patients with osteoarthritis of the knee: an overview of systematic reviews. Jamtvedt G, Dahm KT, Christie A, et al. 2008;88:123-136.

Use of joint mobilization in a patient with severely restricted hip motion following bilateral hip resurfacing arthroplasty. Crow JB, Gelfand B, Su EP. 2008;88:1591-1600.

Rheumatoid Arthritis

Keitel Functional Test for patients with rheumatoid arthritis: translation, reliability, validity, and responsiveness. Holm B, Jacobsen S, Skjodt H, et al. 2008;88:664-678.

Views on exercise maintenance: variations among patients with rheumatoid arthritis. Swärdh E, Biguet G, Opava CH. 2008;88:1049-1060.

- Correction. 2008;88:1232.

B

Balance

Balance and eye movement training to improve gait in people with progressive supranuclear palsy: a quasi-randomized clinical trial. Zampieri C, Di Fabio RP. 2008;88:1460-1473.

Diagnosis of fall risk in Parkinson disease: an analysis of individual and collective clinical balance test interpretation. Dibble LE, Christensen J, Ballard DJ, Foreman KB. 2008;88:323-332.

- Letter to the editor. Norton BJ. 2008;88:679.
- Author response. Dibble LE, Foreman KB, Ballard DJ, Christensen J. 2008;88:679-680.

Impact of health perception, balance perception, fall history, balance performance, and gait speed on walking activity in older adults. Talkowski JB, Brach JS, Studenski S, Newman AB. 2008;88:1474-1481.

Increased risk of falling in older community-dwelling women with mild cognitive impairment. Liu-Ambrose TY, Ashe MC, Graf P, et al. 2008;88:1482-1491.

Temporal and spatial characteristics of gait during performance of the Dynamic Gait Index in people with and people without balance or vestibular disorders. Marchetti GF, Whitney SL, Blatt PJ, et al. 2008;88:640-651.

Use of the Berg Balance Scale for predicting multiple falls in community-dwelling elderly people: a prospective study. Muir SW, Berg K, Chesworth B, Speechley M. 2008;88:449-459.

- Invited commentary. Studenski SA. 2008;88:460-461.
- Correction. 2008;88:688.

Biofeedback

Influence of feedback schedule in motor performance and learning of a lumbar multifidus muscle task using rehabilitative ultrasound imaging: a randomized clinical trial. Herbert WJ, Heiss DG, Basso DM. 2008;88:261-269.

C

Cardiac

General

A four-week, task-specific neuroprosthesis program for a

person with no active wrist or finger movement because of chronic stroke. Dunning K, Berberich A, Albers B, et al. 2008;88:397–405.

- Letter to the editor. Broetz D, Soekadar SR, Birbaumer N. 2008;88:970.
- Author response. Dunning K, Hill Hermann V, Levine PG. 2008;88:970–972.

Group-based aerobic interval training in patients with chronic heart failure: Norwegian Ullevaal Model. Nilsson BB, Hellesnes B, Westheim A, Risberg MA. 2008;88:523–535.

Interrater reliability of the Extended ICF Core Set for stroke applied by physical therapists. Starrost K, Geyh S, Trautwein A, et al. 2008;88:841–851.

- Invited commentary. Jette AM. 2008;88:851–853.
- Author response. Geyh S, Stucki G, Cieza A. 2008;88:854–856.

Modified constraint-induced therapy in chronic stroke: results of a single-blinded randomized controlled trial. Page SJ, Levine P, Leonard A, et al. 2008;88:333–340.

- Letter to the editor. Wolf SL. 2008;88:680–684.
- Author response. Page S, Levine P. 2008;88:684–688.

Reproducibility and minimal detectable change of three-dimensional kinematic analysis of reaching tasks in people with hemiparesis after stroke. Wagner JM, Rhodes JA, Patten C. 2008;88:652–663.

Running training after stroke: a single-subject report. Miller EW, Combs SA, Fish C, et al. 2008;88:511–522.

Scale for Contraversive Pushing: cutoff scores for diagnosing "pusher behavior" and construct validity. Baccini M, Paci M, Nannetti L, et al. 2008;88:947–955.

Therapeutic effect of an implantable peroneal nerve stimulator in subjects with chronic stroke and footdrop: a randomized controlled trial. Kottink AIR, Hermens HJ, Nene AV, et al. 2008;88:437–448.

Usefulness of the Berg Balance Scale in stroke rehabilitation: a systematic review. Blum L, Korner-Bitensky N. 2008;88:559–566.

Tests and Measurements

Cardiorespiratory capacity after weight-supported treadmill training in patients with traumatic brain injury. Mossberg KA, Orlander EE, Norcross JL. 2008;88:77–87.

Hierarchical properties of the motor function sections of the Fugl-Meyer Assessment Scale for people after stroke: a retrospective study. Crow JL, Harmeling-van der Wel BC. 2008;88:1554–1567.

Modified total-body recumbent stepper exercise test for assessing peak oxygen consumption in people with chronic stroke. Billinger SA, Tseng BY, Kluding PM. 2008;88:1188–1195.

Cerebral Palsy

Evaluation

Reference curves for the Gross Motor Function Measure: percentiles for clinical description and tracking over time among children with cerebral palsy. Hanna SE, Bartlett DJ, Rivard LM, Russell DJ. 2008;88:596–607.

Roles of reflex activity and co-contraction during assessments of spasticity of the knee flexor and knee extensor muscles in children with cerebral palsy and different functional levels. Pierce SR, Barbe MF, Barr AE, et al. 2008;88:1124–1134.

Treatment

Use of a low-cost, commercially available gaming console (Wii) for rehabilitation of an adolescent with cerebral palsy. Deutsch JE, Borbely M, Filler J, et al. 2008;88:1196–1207.

Computers

Computerized visual feedback: an adjunct to robotic-assisted gait training. Banz R, Bolliger M, Colombo G, et al. 2008;88:1135–1145.

Gait variability in older adults: observational rating validated by comparison with a computerized walkway gold standard. Huang WNW, VanSwearingen JM, Brach JS. 2008;88:1146–1153.

D

Diabetes

Clinical measures of physical fitness predict insulin resistance in people at risk for diabetes. Chen CN, Chuang LM, Wu YT. 2008;88:1355–1364.

Comparison of combined aerobic and high-force eccentric resistance exercise with aerobic exercise only for people with type 2 diabetes mellitus. Marcus RL, Smith S, Morrell G, et al. 2008;88:1345–1354.

Diabetes and associated risk factors in patients referred for physical therapy in a national primary care electronic medical record database. Kirkness CS, Marcus RL, LaStayo PC, et al. 2008;88:1408–1416.

Diabetes mellitus and gait dysfunction: possible explanatory factors. Brach JS, Talkowski JB, Strotmeyer ES, Newman AB. 2008;88:1365–1374.

Diabetes-related microvascular and macrovascular diseases in the physical therapy setting. Cade WT. 2008;88:1322–1335.

Economic realities associated with diabetes care: opportunities to expand delivery of physical therapist services to a vulnerable population. Cohn R. 2008;88:1417–1424.

Editorial: People with diabetes: a population desperate for movement. Mueller MJ. 2008;88:1250–1253.

Effect of weight-bearing activity on foot ulcer incidence in people with diabetic peripheral neuropathy: Feet First randomized controlled trial. LeMaster JW, Mueller MJ, Reiber GE, et al. 2008;88:1385–1398.

Epidemiology of diabetes and diabetes-related complications. Deshpande AD, Harris-Hayes M, Schootman M. 2008;88:1254–1264.

Excessive adipose tissue infiltration in skeletal muscle in individuals with obesity, diabetes mellitus, and peripheral neuropathy: association

with performance and function. Hilton TN, Tuttle LJ, Bohnert KL, et al. 2008;88:1336-1344.

Exercise and glycemic control in diabetes: benefits, challenges, and adjustments to pharmacotherapy. Gulve EA. 2008;88:1297-1321.

Inflammatory osteolysis in diabetic neuropathic (Charcot) arthropathies of the foot. Sinacore DR, Hastings MK, Bohnert KL, et al. 2008;88:1399-1407.

Intricacies of fat. Stehno-Bittel L. 2008;88:1265-1278.

Physical activity and diabetes: opportunities for prevention through policy. Deshpande AD, Dodson EA, Gorman I, Brownson RC. 2008;88:1425-1435.

Plantar stresses on the neuropathic foot during barefoot walking. Mueller MJ, Zou D, Bohnert KL, et al. 2008;88:1375-1384.

Reprint—Comprehensive foot examination and risk assessment: a report of the Foot Care Interest Group of the American Diabetes Association, with endorsement by the American Association of Clinical Endocrinologists. [Reprinted from *Diabetes Care*. 2008;31:1679-1685.] Boulton AJM, Armstrong DG, Albert SF, et al. 2008;88:1436-1443.

Skeletal muscle insulin resistance: roles of fatty acid metabolism and exercise. Turcotte LP, Fisher JS. 2008;88:1279-1296.

Diagnosis

Misdiagnosis of serotonin syndrome as fibromyalgia and the role of physical therapists. Alnwick GM. 2008;88:757-765.

Process for applying the *International Classification of Functioning, Disability and Health* model to a patient with patellar dislocation. Helgeson K, Smith AR Jr. 2008;88:956-964.

Scale for Contraversive Pushing: cutoff scores for diagnosing "pusher behavior" and construct validity. Baccini M, Paci M, Nannetti L, et al. 2008;88:947-955.

E

Editorials

Editorial: Climbing out of our silos to improve practice. Craik RL. 2008;88:555-558.

Editorial: Does clinical education need a *series* of tools to assess success? Craik RL. 2008;88:1106-1108.

Editorial: In memoriam: Beverly Petterson Bishop (1922-2008). Craik RL, Harris S, Helders PJ. 2008;88:1454-1456.

Editorial: New author service: PTJ makes PubMed Central deposits. Editorial Office. 2008;88:699-700.

Editorial: People with diabetes: a population desperate for movement. Mueller MJ. 2008;88:1250-1253.

Editorial: Still "not satisfied"... yet. Craik RL, Maher C. 2008;88:423-425.

- Correction. 2008;88:688.

Editorial: The 2008/2009 editorial board. Craik RL. 2008;88:986-988.

Guest editorial: "What's in a name?" revisited. Behrman AL, Plummer-D'Amato P. 2008;88:6-9.

Education: Patient/Family

Effects of education on return-to-work status for people with fear-avoidance beliefs and acute low back pain. Godges JJ, Anger MA, Zimmerman G, Delitto A. 2008;88:231-239.

Education: Physical Therapist

Clinical Education

Editorial: Does clinical education need a *series* of tools to assess success? Craik RL. 2008;88:1106-1108.

Pediatric physical therapists' perceptions of their training in assistive technology. Long TM, Perry DF. 2008;88:629-639.

Electromyography

Electromyography for assessment of pain in low back muscles. Candotti CT, Loss JF, Pressi AMS, et al. 2008;88:1061-1067.

Electrotherapy

Electrical Stimulation

Effect of burst frequency and duration of kilohertz-frequency alternating currents and of low-frequency pulsed currents on strength of contraction, muscle fatigue, and perceived discomfort. Laufer Y, Elboim M. 2008;88:1167-1176.

Therapeutic effect of an implantable peroneal nerve stimulator in subjects with chronic stroke and footdrop: a randomized controlled trial. Kottink AIR, Hermens HJ, Nene AV, et al. 2008;88:437-448.

Using customized rate-coding and recruitment strategies to maintain forces during repetitive activation of human muscles. Chou LW, Kesar TM, Binder-Macleod SA. 2008;88:363-375.

Iontophoresis

In vitro optimization of dexamethasone phosphate delivery by iontophoresis. Sylvestre JP, Guy RH, Delgado-Charro MB. 2008;88:1177-1185.

- Invited commentary. Underwood FB. 2008;88:1185-1187.

Equipment

General

Use of a low-cost, commercially available gaming console (Wii) for rehabilitation of an adolescent with cerebral palsy. Deutsch JE, Borbely M, Filler J, et al. 2008;88:1196-1207.

Evidence-Based Practice

A description of the trials, reviews, and practice guidelines indexed in the PEDro database. Maher CG, Moseley AM, Sherrington C, et al. 2008;88:1068-1077.

Ottawa Panel Evidence-Based Clinical Practice Guidelines for Aerobic Fitness Exercises in the Management of Fibromyalgia: Part 1. Brosseau L, Wells GA, Tugwell P, et al. 2008;88:857-871.

Ottawa Panel Evidence-Based Clinical Practice Guidelines for Strengthening Exercises in the Management of Fibromyalgia: Part 2. Brosseau L, Wells GA, Tugwell P, et al. 2008;88:873-886.

Exercise

General

Cardiorespiratory capacity after weight-supported treadmill training in patients with traumatic brain injury. Mossberg KA, Orlander EE, Norcross JL. 2008;88:77-87.

Comparison of combined aerobic and high-force eccentric resistance exercise with aerobic exercise only for people with type 2 diabetes mellitus. Marcus RL, Smith S, Morrell G, et al. 2008;88:1345-1354.

Effects of a single session of posterior-to-anterior spinal mobilization and press-up exercise on pain response and lumbar spine extension in people with nonspecific low back pain. Powers CM, Beneck GJ, Kulig K, et al. 2008;88:485-493.

Effects of intensity of treadmill training on developmental outcomes and stepping in infants with Down syndrome: a randomized trial. Ulrich DA, Lloyd MC, Tiernan CW, et al. 2008;88:114-122.

Effects of Pilates exercises on shoulder range of motion, pain, mood, and upper-extremity function in women living with breast cancer: a pilot study. Keays KS, Harris SR, Lucyshyn JM, MacIntyre DL. 2008;88:494-510.

Effects of whole-body vibration exercise on lower-extremity muscle strength and power in an older population: a randomized clinical trial. Rees SS, Murphy AJ, Watsford ML. 2008;88:462-470.

Endurance exercise training to improve economy of movement of people with Parkinson disease: three case reports. Schenkman M, Hall D, Kumar R, Kohrt WM. 2008;88:63-76.

Exercise and glycemic control in diabetes: benefits, challenges, and adjustments to pharmacotherapy. Gulve EA. 2008;88:1297-1321.

Group-based aerobic interval training in patients with chronic heart failure: Norwegian Ullevaal Model. Nilsson BB, Hellesnes B, Westheim A, Risberg MA. 2008;88:523-535.

Hydrotherapy versus conventional land-based exercise for the management of patients with osteoarthritis of the knee: a randomized clinical trial. Silva LE, Valim V, Pessanha APC, et al. 2008;88:12-21.

Locomotor training using body-weight support on a treadmill in conjunction with ongoing physical therapy in a child with severe cerebellar ataxia. Cernak K, Stevens V, Price R, Shumway-Cook A. 2008;88:88-97.

Modified total-body recumbent stepper exercise test for assessing peak oxygen consumption in people with chronic stroke. Billinger SA, Tseng BY, Kluding PM. 2008;88:1188-1195.

Ottawa Panel Evidence-Based Clinical Practice Guidelines for Aerobic Fitness Exercises in the Management of Fibromyalgia: Part 1. Brosseau L, Wells GA, Tugwell P, et al. 2008;88:857-871.

Physical therapists' use of therapeutic exercise for patients with clinical knee osteoarthritis in the United Kingdom: in line with current recommendations? Holden MA, Nicholls EE, Hay EM, Foster NE. 2008;88:1109-1121.

- Invited commentary. Lund H. 2008;88:1121-1122.
- Author response. Foster NE. 2008;88:1123.

Running training after stroke: a single-subject report. Miller EW, Combs SA, Fish C, et al. 2008;88:511-522.

Skeletal muscle insulin resistance: roles of fatty acid metabolism and exercise. Turcotte LP, Fisher JS. 2008;88:1279-1296.

Validity of values for metabolic equivalents of task during submaximal all-extremity exercise and reliability of exercise responses in frail older adults. Mendelsohn ME, Connelly DM, Overend TJ, Petrella PJ. 2008;88:747-756.

Views on exercise maintenance: variations among patients with rheumatoid arthritis. Swärdh E, Biguet G, Opava CH. 2008;88:1049-1060.

- Correction. 2008;88:1232.

Strengthening

Continence and quality-of-life outcomes 6 months following an intensive pelvic-floor muscle exercise program for female stress urinary incontinence: a randomized trial comparing low- and high-frequency maintenance exercise. Borello-France DF, Downey PA, Zyczynski HM, Rause CR. 2008;88:1545-1553.

Investigation of clinical effects of high- and low-resistance training for patients with knee osteoarthritis: a randomized controlled trial. Jan MH, Lin JJ, Liau JJ, et al. 2008;88:427-436.

Ottawa Panel Evidence-Based Clinical Practice Guidelines for Strengthening Exercises in the Management of Fibromyalgia: Part 2. Brosseau L, Wells GA, Tugwell P, et al. 2008;88:873-886.

Use of transabdominal ultrasound imaging in retraining the pelvic-floor muscles of a woman postpartum. Ariail A, Sears T, Hampton E. 2008;88:1208-1217.

F

Fatigue

Effect of burst frequency and duration of kilohertz-frequency alternating currents and of low-frequency pulsed currents on strength of contraction, muscle fatigue, and perceived discomfort. Laufer Y, Elboim M. 2008;88:1167-1176.

Foundation for Physical Therapy

Scholarships, Fellowships, and Grants. 2008;88:140-141, 297-298, 413-414, 543-544, 689-690, 801-802, 895-896, 975-976, 1097-1098, 1235-1236, 1444-1445, 1607-1608.

Functional Training and Activities

Effect of weight-bearing activity on foot ulcer incidence in people with diabetic peripheral neuropathy: Feet First randomized controlled trial. LeMaster JW, Mueller MJ, Reiber GE, et al. 2008;88:1385-1398.

Habitual physical activity behavior of patients after primary total hip arthroplasty. Wagenmakers R, Stevens M, Zijlstra W, et al. 2008;88:1039-1048.

Physical activity and diabetes: opportunities for prevention through policy. Deshpande AD, Dodson EA, Gorman I, Brownson RC. 2008;88:1425–1435.

G

Geriatrics

Associations between physical examination and self-reported physical function in older community-dwelling adults with knee pain. Wood L, Peat G, Thomas E, et al. 2008;88:33–42.

Impact of health perception, balance perception, fall history, balance performance, and gait speed on walking activity in older adults. Talkowski JB, Brach JS, Studenski S, Newman AB. 2008;88:1474–1481.

Increased risk of falling in older community-dwelling women with mild cognitive impairment. Liu-Ambrose TY, Ashe MC, Graf P, et al. 2008;88:1482–1491.

Validity of values for metabolic equivalents of task during submaximal all-extremity exercise and reliability of exercise responses in frail older adults. Mendelsohn ME, Connelly DM, Overend TJ, Petrella PJ. 2008;88:747–756.

H

Head Injury

Cardiorespiratory capacity after weight-supported treadmill training in patients with traumatic brain injury. Mossberg KA, Orlander EE, Norcross JL. 2008;88:77–87.

Time series analysis of spontaneous upper-extremity movements of premature infants with brain injuries. Ohgi S, Morita S, Loo KK, Mizuike C. 2008;88:1022–1033.

- Invited commentary. Fetters L, Scholz JP. 2008;88:1034–1036.
- Author response. Ohgi S, Morita S, Loo KK, Mizuike C. 2008;88:1037–1038.

Health Care

Health care providers' knowledge, attitudes, and self-efficacy for working with patients with spinal cord injury who have diverse sexual orientations. Burch A. 2008;88:191–198.

Implementing an integrated electronic outcomes and electronic health record process to create a foundation for clinical practice improvement. Deutscher D, Hart DL, Dickstein R, et al. 2008;88:270–285.

Hemiplegia

General

Reproducibility and minimal detectable change of three-dimensional kinematic analysis of reaching tasks in people with hemiparesis after stroke. Wagner JM, Rhodes JA, Patten C. 2008;88:652–663.

Orthoses and Supports

A four-week, task-specific neuroprosthesis program for a person with no active wrist or finger movement because of chronic stroke. Dunning K, Berberich A, Albers B, et al. 2008;88:397–405.

- Letter to the editor. Broetz D, Soekadar SR, Birbaumer N. 2008;88:970.
- Author response. Dunning K, Hill Hermann V, Levine PG. 2008;88:970–972.

Home Care

Interventions used by physical therapists in home care for people after hip fracture. Mangione KK, Lopopolo RB, Neff NP, et al. 2008;88:199–210.

Hydrotherapy

Hydrotherapy versus conventional land-based exercise for the management of patients with osteoarthritis of the knee: a randomized clinical trial. Silva LE, Valim V, Pessanha APC, et al. 2008;88:12–21.

K

Kinesiology/Biomechanics

Gait Analysis

Balance and eye movement training to improve gait in people with progressive supranuclear palsy: a quasi-randomized clinical trial. Zampieri C, Di Fabio RP. 2008;88:1460–1473.

Computerized visual feedback: an adjunct to robotic-assisted gait training. Banz R, Bolliger M, Colombo G, et al. 2008;88:1135–1145.

Diabetes mellitus and gait dysfunction: possible explanatory factors. Brach JS, Talkowski JB, Strotmeyer ES, Newman AB. 2008;88:1365–1374.

Gait characteristics of a soldier with a traumatic hip disarticulation. Schnall BL, Baum BS, Andrews AM. 2008;88:1568–1577.

Gait variability in older adults: observational rating validated by comparison with a computerized walkway gold standard. Huang WNW, VanSwearingen JM, Brach JS. 2008;88:1146–1153.

Impact of health perception, balance perception, fall history, balance performance, and gait speed on walking activity in older adults. Talkowski JB, Brach JS, Studenski S, Newman AB. 2008;88:1474–1481.

Plantar stresses on the neuropathic foot during barefoot walking. Mueller MJ, Zou D, Bohnert KL, et al. 2008;88:1375–1384.

Running training after stroke: a single-subject report. Miller EW, Combs SA, Fish C, et al. 2008;88:511–522.

Temporal and spatial characteristics of gait during performance of the Dynamic Gait Index in people with and people without balance or vestibular disorders. Marchetti GF, Whitney SL, Blatt PJ, et al. 2008;88:640–651.

General

Ilial anterior rotation hypermobility in a female collegiate tennis player. Vaughn HT, Nitsch W. 2008;88:1578–1590.

Persistence of altered movement patterns during a sit-to-stand task 1 year following unilateral total knee arthroplasty. Farquhar SJ, Reisman DS, Snyder-Mackler L. 2008;88:567–579.

- Correction. 2008;88:797.

Reproducibility and minimal detectable change of three-dimensional kinematic analysis of reaching tasks in people with hemiparesis after stroke. Wagner JM, Rhodes JA, Patten C. 2008;88:652-663.

Lower Extremity

Influence of different footwear on force of landing during running. Cheung RTH, Ng GYF. 2008;88:620-628.

- Correction. 2008;88:797.

L

Locomotion

Locomotor training restores walking in a nonambulatory child with chronic, severe, incomplete cervical spinal cord injury. Behrman AL, Nair PM, Bowden MG, et al. 2008;88:580-590.

- Invited commentary. Edgerton VR. 2008;88:590-591.
- Invited commentary. McEwen IR. 2008;88:592-594.
- Author response. Behrman AL, Nair PM, Bowden MG, et al. 2008;88:594-595.

Locomotor training using body-weight support on a treadmill in conjunction with ongoing physical therapy in a child with severe cerebellar ataxia. Cernak K, Stevens V, Price R, Shumway-Cook A. 2008;88:88-97.

Lower Extremity

Ankle and Foot

Bone mineral density of the tarsals and metatarsals with reloading. Hastings MK, Gelber J, Commean PK, et al. 2008;88:766-779.

Effect of weight-bearing activity on foot ulcer incidence in people with diabetic peripheral neuropathy: Feet First randomized controlled trial. LeMaster JW, Mueller MJ, Reiber GE, et al. 2008;88:1385-1398.

Gaze control and foot kinematics during stair climbing: characteristics leading to fall risk in progressive supranuclear palsy. Di Fabio RP, Zampieri C, Tuite P. 2008;88:240-250.

Inflammatory osteolysis in diabetic neuropathic (Charcot) arthropathies of the foot. Sinacore DR, Hastings MK, Bohnert KL, et al. 2008;88:1399-1407.

Plantar stresses on the neuropathic foot during barefoot walking. Mueller MJ, Zou D, Bohnert KL, et al. 2008;88:1375-1384.

Reprint—Comprehensive foot examination and risk assessment: a report of the Foot Care Interest Group of the American Diabetes Association, with endorsement by the American Association of Clinical Endocrinologists. [Reprinted from *Diabetes Care*. 2008;31:1679-1685.] Boulton AJM, Armstrong DG, Albert SF, et al. 2008;88:1436-1443.

General

Estimation of thigh muscle mass with magnetic resonance imaging in older adults and people with chronic obstructive pulmonary disease. Mathur S, Takai KPR, MacIntyre DL, Reid D. 2008;88:219-230.

Hip

[Effectiveness of nonpharmacological and nonsurgical interventions for hip osteoarthritis: an umbrella review of high-quality systematic reviews. Christie A, Jamtvedt G, Dahm KT, et al. 2007;87:1716-1727.]

- Letter to the editor. Leeb BF, Rintelen B. 2008;88:407-408.
- Author response. Moe RH, Haavardsholm EA, Christie A, et al. 2008;88:408.

Functional outcomes and quality of life after tumor-related hemipelvectomy. Beck LA, Einertson MJ, Winemiller MH, et al. 2008;88:916-927.

Gait characteristics of a soldier with a traumatic hip disarticulation. Schnall BL, Baum BS, Andrews AM. 2008;88:1568-1577.

Habitual physical activity behavior of patients after primary total hip arthroplasty. Wagenmakers R, Stevens M, Zijlstra W, et al. 2008;88:1039-1048.

Interventions used by physical therapists in home care for people after hip fracture. Mangione KK, Lopopolo RB, Neff NP, et al. 2008;88:199-210.

Physical functioning before and after total hip arthroplasty: perception and performance. van den Akker-Scheek I, Zijlstra W, Groothoff JW, et al. 2008;88:712-719.

Predictive value of the Western Ontario and McMaster Universities Osteoarthritis Index for the amount of physical activity after total hip arthroplasty. Wagenmakers R, Stevens M, van den Akker-Scheek I, et al. 2008;88:211-218.

Traumatic dislocation of the hip in a high school football player. Yates C, Bandy WD, Blasier RD. 2008;88:780-788.

Use of joint mobilization in a patient with severely restricted hip motion following bilateral hip resurfacing arthroplasty. Crow JB, Gelfand B, Su EP. 2008;88:1591-1600.

Knee

After partial knee replacement, patients can kneel, but they need to be taught to do so: a single-blind randomized controlled trial. Jenkins C, Barker KL, Pandit H, et al. 2008;88:1012-1021.

Assessing recovery and establishing prognosis following total knee arthroplasty. Kennedy DM, Stratford PW, Riddle DL, et al. 2008;88:22-32.

Associations between physical examination and self-reported physical function in older community-dwelling adults with knee pain. Wood L, Peat G, Thomas E, et al. 2008;88:33-42.

Hydrotherapy versus conventional land-based exercise for the management of patients with osteoarthritis of the knee: a randomized clinical trial. Silva LE, Valim V, Pessanha APC, et al. 2008;88:12-21.

Instability, laxity, and physical function in patients with medial knee osteoarthritis. Schmitt LC, Fitzgerald GK, Reisman AS, Rudolph KS. 2008;88:1506-1516.

Investigation of clinical effects of high- and low-resistance training for patients with knee osteoarthritis: a randomized controlled trial. Jan MH, Lin JJ, Liau JJ, et al. 2008;88:427-436.

Persistence of altered movement patterns during a sit-to-stand task 1 year following unilateral total knee arthroplasty. Farquhar SJ, Reisman DS, Snyder-Mackler L. 2008;88: 567-579.

- Correction. 2008;88:797.

Physical therapists' use of therapeutic exercise for patients with clinical knee osteoarthritis in the United Kingdom: in line with current recommendations? Holden MA, Nicholls EE, Hay EM, Foster NE. 2008;88:1109-1121.

- Invited commentary. Lund H. 2008;88:1121-1122.
- Author response. Foster NE. 2008;88:1123.

Physical therapy interventions for patients with osteoarthritis of the knee: an overview of systematic reviews. Jamtvedt G, Dahm KT, Christie A, et al. 2008;88:123-136.

Process for applying the *International Classification of Functioning, Disability and Health* model to a patient with patellar dislocation. Helgeson K, Smith AR Jr. 2008;88:956-964.

Roles of reflex activity and co-contraction during assessments of spasticity of the knee flexor and knee extensor muscles in children with cerebral palsy and different functional levels. Pierce SR, Barbe MF, Barr AE, et al. 2008;88: 1124-1134.

M

Medical Conditions

Relationships among severity of osteonecrosis, pain, range of motion, and functional mobility in children, adolescents, and young adults with acute lymphoblastic leukemia. Marchese VG, Connolly BH, Able C, et al. 2008;88:341-350.

Motor Ability

Reproducibility and minimal detectable change of three-dimensional kinematic analysis of reaching tasks in people with hemiparesis after stroke. Wagner JM, Rhodes JA, Patten C. 2008;88: 652-663.

Motor Activity

Influence of feedback schedule in motor performance and learning of a lumbar multifidus muscle task using rehabilitative ultrasound imaging: a randomized clinical trial. Herbert WJ, Heiss DG, Basso DM. 2008;88:261-269.

Motor Skills

Effect of focus of attention and age on motor acquisition, retention, and transfer: a randomized trial. Emanuel M, Jarus T, Bart O. 2008;88:251-260.

Modified constraint-induced therapy in chronic stroke: results of a single-blinded randomized controlled trial. Page SJ, Levine P, Leonard A, et al. 2008;88:333-340.

- Letter to the editor. Wolf SL. 2008;88:680-684.
- Author response. Page S, Levine P. 2008;88:684-688.

Motor learning in children: feedback effects on skill acquisition. Sullivan KJ, Kantak SS, Burtner PA. 2008;88:720-732.

Multidimensional motor sequence learning is impaired in older but not younger or middle-aged adults. Boyd LA, Vidoni ED, Siengsukon CF. 2008;88:351-362.

Movement Science

Endurance exercise training to improve economy of movement of people with Parkinson disease: three case reports. Schenkman M, Hall D, Kumar R, Kohrt WM. 2008;88:63-76.

Gaze control and foot kinematics during stair climbing: characteristics leading to fall risk in progressive supranuclear palsy. Di Fabio RP, Zampieri C, Tuite P. 2008;88: 240-250.

Ilial anterior rotation hypermobility in a female collegiate tennis player. Vaughn HT, Nitsch W. 2008;88: 1578-1590.

Influence of different footwear on force of landing during running. Cheung RTH, Ng GYF. 2008;88: 620-628.

- Correction. 2008;88:797.

Locomotor training restores walking in a nonambulatory child with chronic, severe, incomplete cervical spinal cord injury. Behrman AL, Nair PM, Bowden MG, et al. 2008;88:580-590.

- Invited commentary. Edgerton VR. 2008;88:590-591.
- Invited commentary. McEwen IR. 2008;88:592-594.
- Author response. Behrman AL, Nair PM, Bowden MG, et al. 2008;88:594-595.

Locomotor training using body-weight support on a treadmill in conjunction with ongoing physical therapy in a child with severe cerebellar ataxia. Cernak K, Stevens V, Price R, Shumway-Cook A. 2008;88:88-97.

Movement training advances the emergence of reaching in infants born at less than 33 weeks of gestational age: a randomized clinical trial. Heathcock JC, Lobo M, Galloway JC. 2008;88:310-322.

Multidimensional motor sequence learning is impaired in older but not younger or middle-aged adults. Boyd LA, Vidoni ED, Siengsukon CF. 2008;88:351-362.

Persistence of altered movement patterns during a sit-to-stand task 1 year following unilateral total knee arthroplasty. Farquhar SJ, Reisman DS, Snyder-Mackler L. 2008;88: 567-579.

- Correction. 2008;88:797.

Puckering and blowing facial expression in people with facial movement disorders. Denlinger RL, VanSwearingen JM, Cohn JF, Schmidt KL. 2008;88:909-915.

- Letter to the editor. Halili A. 2008;88:1231.
- Author response. Schmidt KL, Denlinger RL, VanSwearingen JM, Cohn JF. 2008;88:1231-1232.

Time series analysis of spontaneous upper-extremity movements of premature infants with brain injuries. Ohgi S, Morita S, Loo KK, Mizuike C. 2008;88:1022-1033.

- Invited commentary. Fetters L, Scholz JP. 2008;88:1034-1036.
- Author response. Ohgi S, Morita S, Loo KK, Mizuike C. 2008;88: 1037-1038.

Muscle

Continence and quality-of-life outcomes 6 months following an intensive pelvic-floor muscle exercise program for female stress urinary incontinence: a randomized trial comparing low- and high-frequency maintenance exercise. Borello-France DF, Downey PA, Zyczynski HM, Rause CR. 2008;88:1545–1553.

Excessive adipose tissue infiltration in skeletal muscle in individuals with obesity, diabetes mellitus, and peripheral neuropathy: association with performance and function. Hilton TN, Tuttle LJ, Bohnert KL, et al. 2008;88:1336–1344.

Ottawa Panel Evidence-Based Clinical Practice Guidelines for Aerobic Fitness Exercises in the Management of Fibromyalgia: Part 1. Brosseau L, Wells GA, Tugwell P, et al. 2008;88:857–871.

Ottawa Panel Evidence-Based Clinical Practice Guidelines for Strengthening Exercises in the Management of Fibromyalgia: Part 2. Brosseau L, Wells GA, Tugwell P, et al. 2008;88:873–886.

Skeletal muscle insulin resistance: roles of fatty acid metabolism and exercise. Turcotte LP, Fisher JS. 2008;88:1279–1296.

Use of transabdominal ultrasound imaging in retraining the pelvic-floor muscles of a woman postpartum. Ariail A, Sears T, Hampton E. 2008;88:1208–1217.

Muscle Performance

General

Effect of burst frequency and duration of kilohertz-frequency alternating currents and of low-frequency pulsed currents on strength of contraction, muscle fatigue, and perceived discomfort. Laufer Y, Elboim M. 2008;88:1167–1176.

Influence of feedback schedule in motor performance and learning of a lumbar multifidus muscle task using rehabilitative ultrasound imaging: a randomized clinical trial. Herbert WJ, Heiss DG, Basso DM. 2008;88:261–269.

Muscle activation during selected strength exercises in women with chronic neck muscle pain. Andersen LL, Kjær M, Andersen CH, et al. 2008;88:703–711.

Using customized rate-coding and recruitment strategies to maintain forces during repetitive activation of human muscles. Chou LW, Kesar TM, Binder-Macleod SA. 2008;88:363–375.

Lower Extremity

Effects of whole-body vibration exercise on lower-extremity muscle strength and power in an older population: a randomized clinical trial. Rees SS, Murphy AJ, Watsford ML. 2008;88:462–470.

Roles of reflex activity and co-contraction during assessments of spasticity of the knee flexor and knee extensor muscles in children with cerebral palsy and different functional levels. Pierce SR, Barbe MF, Barr AE, et al. 2008;88:1124–1134.

Therapeutic effect of an implantable peroneal nerve stimulator in subjects with chronic stroke and footdrop: a randomized controlled trial. Kottink AIR, Hermens HJ, Nene AV, et al. 2008;88:437–448.

Muscular Dystrophy

Facioscapulohumeral dystrophy. Pandya S, King WM, Tawil R. 2008;88:105–113.

Musculoskeletal Conditions

Misdiagnosis of serotonin syndrome as fibromyalgia and the role of physical therapists. Alnwick GM. 2008;88:757–765.

Work-related musculoskeletal disorders in physical therapists: a prospective cohort study with 1-year follow-up. Campo M, Weiser S, Koenig KL, Nordin M. 2008;88:608–619.

- Letter to the editor. Halloran SW. 2008;88:887.
- Author response. Campo MA. 2008;88:887–888.

N

Neck and Trunk

Back

[A comparison of a modified Oswestry Low Back Pain Disability Questionnaire and the Quebec Back Pain Disability Scale. Fritz JM, Irrgang JJ. 2001;81:776–788.]

- Correction. 2008;88:138–139.

Effects of a single session of posterior-to-anterior spinal mobilization and press-up exercise on pain response and lumbar spine extension in people with nonspecific low back pain. Powers CM, Beneck GJ, Kulig K, et al. 2008;88:485–493.

Effects of education on return-to-work status for people with fear-avoidance beliefs and acute low back pain. Godges JJ, Anger MA, Zimmerman G, Delitto A. 2008;88:231–239.

Electromyography for assessment of pain in low back muscles. Candotti CT, Loss JF, Pressi AMS, et al. 2008;88:1061–1067.

Factors influencing the use of outcome measures for patients with low back pain: a survey of New Zealand physical therapists. Copeland JM, Taylor WJ, Dean SG. 2008;88:1492–1505.

[Indicators of lumbar zygapophyseal joint pain: survey of an expert panel with the Delphi technique. Wilde VE, Ford JJ, McMeeken JM. 2007;87:1348–1361.]

- Letter to the editor. Strobl RM. 2008;88:137.
- Author response. Wilde VE, McMeeken JM. 2008;88:137.

Predictors of physical therapy clinic performance in the treatment of patients with low back pain syndromes. Resnik L, Liu D, Mor V, Hart DL. 2008;88:989–1004.

- Invited commentary. Freburger JK, Mielenz TJ. 2008;88:1004–1006.
- Invited commentary. Fritz JM. 2008;88:1007–1009.
- Author response. Resnik L, Hart DL. 2008;88:1009–1011.

General

Intertester reliability and validity of motion assessments during lumbar spine accessory motion testing. Landel R, Kulig K, Fredericson M, et al. 2008;88:43–49.

- Letter to the editor. Abbott JH. 2008;88:536.
- Author response. Landel R, Kulig K, Fredericson M, et al. 2008;88:536–537.

Undetected hangman's fracture in a patient referred for physical therapy for the treatment of neck pain following trauma. Ross MD, Cheeks JM. 2008;88:98–104.

Neck

Muscle activation during selected strength exercises in women with chronic neck muscle pain. Andersen LL, Kjær M, Andersen CH, et al. 2008;88:703–711.

Neuromuscular Disorders

General

Excessive adipose tissue infiltration in skeletal muscle in individuals with obesity, diabetes mellitus, and peripheral neuropathy: association with performance and function. Hilton TN, Tuttle LJ, Bohnert KL, et al. 2008;88:1336–1344.

Gaze control and foot kinematics during stair climbing: characteristics leading to fall risk in progressive supranuclear palsy. Di Fabio RP, Zampieri C, Tuite P. 2008;88:240–250.

Locomotor training using body-weight support on a treadmill in conjunction with ongoing physical therapy in a child with severe cerebellar ataxia. Cernak K, Stevens V, Price R, Shumway-Cook A. 2008;88:88–97.

Physical therapy intervention for people with Huntington disease. Busse ME, Khalil H, Quinn L, Rosser AE. 2008;88:820–831.

Multiple Sclerosis

Measuring deterioration in International Classification of Functioning domains of people with multiple sclerosis who are ambulatory. Paltamaa J, Sarasoja T, Leskinen E, et al. 2008;88:176–190.

Parkinson Disease

Diagnosis of fall risk in Parkinson disease: an analysis of individual and collective clinical balance test interpretation. Dibble LE, Christensen J, Ballard DJ, Foreman KB. 2008;88:323–332.

- Letter to the editor. Norton BJ. 2008;88:679.
- Author response. Dibble LE, Foreman KB, Ballard DJ, Christensen J. 2008;88:679–680.

Effectiveness of an inpatient multidisciplinary rehabilitation program for people with Parkinson disease. Ellis T, Katz DI, White DK, et al. 2008;88:812–819.

Endurance exercise training to improve economy of movement of people with Parkinson disease: three case reports. Schenkman M, Hall D, Kumar R, Kohrt WM. 2008;88:63–76.

Test-retest reliability and minimal detectable change on balance and ambulation tests, the 36-Item Short-Form Health Survey, and the Unified Parkinson Disease Rating Scale in people with parkinsonism. Steffen T, Seney M. 2008;88:733–746.

- Letter to the editor. Stratford PW. 2008;88:888–890.
- Author response. Steffen TM, Seney M. 2008;88:891.

O

Oncology

Effects of Pilates exercises on shoulder range of motion, pain, mood, and upper-extremity function in women living with breast cancer: a pilot study. Keays KS, Harris SR, Lucyshyn JM, MacIntyre DL. 2008;88:494–510.

Orthopedics

Fractures

Interventions used by physical therapists in home care for people after hip fracture. Mangione KK, Lopopolo RB, Neff NP, et al. 2008;88:199–210.

Undetected hangman's fracture in a patient referred for physical therapy for the treatment of neck pain following trauma. Ross MD, Cheeks JM. 2008;88:98–104.

General

Assessing recovery and establishing prognosis following total knee arthroplasty. Kennedy DM, Stratford PW, Riddle DL, et al. 2008;88:22–32.

Dose estimation and surveillance of mechanical loading interventions for bone loss after spinal cord injury. Dudley-Javoroski S, Shields RK. 2008;88:387–396.

Physical functioning before and after total hip arthroplasty: perception and performance. van den Akker-Scheek I, Zijlstra W, Groothoff JW, et al. 2008;88:712–719.

Relationships among severity of osteonecrosis, pain, range of motion, and functional mobility in children, adolescents, and young adults with acute lymphoblastic leukemia. Marchese VG, Connolly BH, Able C, et al. 2008;88:341–350.

Orthoses/Splints/Casts

Lower Extremity

Therapeutic effect of an implantable peroneal nerve stimulator in subjects with chronic stroke and footdrop: a randomized controlled trial. Kottink AIR, Hermens HJ, Nene AV, et al. 2008;88:437–448.

P

Pain

Associations between physical examination and self-reported physical function in older community-dwelling adults with knee pain. Wood L, Peat G, Thomas E, et al. 2008;88:33–42.

[A comparison of a modified Oswestry Low Back Pain Disability Questionnaire and the Quebec Back Pain Disability Scale. Fritz JM, Irrgang JJ. 2001;81:776–788.]

- Correction. 2008;88:138–139.

Effects of a single session of posterior-to-anterior spinal mobilization and press-up exercise on pain response and lumbar spine extension in people with nonspecific low back pain. Powers CM, Beneck GJ, Kulig K, et al. 2008;88:485–493.

Effects of education on return-to-work status for people with fear-avoidance beliefs and acute low back pain. Godges JJ, Anger MA, Zimmerman G, Delitto A. 2008;88:231–239.

Electromyography for assessment of pain in low back muscles. Candotti CT, Loss JF, Pressi AMS, et al. 2008;88:1061-1067.

Factors influencing the use of outcome measures for patients with low back pain: a survey of New Zealand physical therapists. Copeland JM, Taylor WJ, Dean SG. 2008;88:1492-1505.

[Indicators of lumbar zygapophyseal joint pain: survey of an expert panel with the Delphi technique. Wilde VE, Ford JJ, McMeeken JM. 2007;87:1348-1361.]

- Letter to the editor. Strobl RM. 2008;88:137.
- Author response. Wilde VE, McMeeken JM. 2008;88:137.

Intertester reliability and validity of motion assessments during lumbar spine accessory motion testing. Landel R, Kulig K, Fredericson M, et al. 2008;88:43-49.

- Letter to the editor. Abbott JH. 2008;88:536.
- Author response. Landel R, Kulig K, Fredericson M, et al. 2008;88:536-537.

Muscle activation during selected strength exercises in women with chronic neck muscle pain. Andersen LL, Kjær M, Andersen CH, et al. 2008;88:703-711.

Predictors of physical therapy clinic performance in the treatment of patients with low back pain syndromes. Resnik L, Liu D, Mor V, Hart DL. 2008;88:989-1004.

- Invited commentary. Freburger JK, Mielenz TJ. 2008:88:1004-1006.
- Invited commentary. Fritz JM. 2008;88:1007-1009.
- Author response. Resnik L, Hart DL. 2008;88:1009-1011.

Undetected hangman's fracture in a patient referred for physical therapy for the treatment of neck pain following trauma. Ross MD, Cheeks JM. 2008;88:98-104.

Pediatrics

Development

Effects of intensity of treadmill training on developmental outcomes and stepping in infants with Down syndrome: a randomized trial. Ulrich DA, Lloyd MC, Tiernan CW, et al. 2008;88:114-122.

Motor learning in children: feedback effects on skill acquisition. Sullivan KJ, Kantak SS, Burtner PA. 2008;88:720-732.

Movement training advances the emergence of reaching in infants born at less than 33 weeks of gestational age: a randomized clinical trial. Heathcock JC, Lobo M, Galloway JC. 2008;88:310-322.

Evaluation

Interrater reliability of the Movement Assessment Battery for Children. Smits-Engelsman BCM, Fiers MJ, Henderson SE, Henderson L. 2008;88:286-294.

Reference curves for the Gross Motor Function Measure: percentiles for clinical description and tracking over time among children with cerebral palsy. Hanna SE, Bartlett DJ, Rivard LM, Russell DJ. 2008;88:596-607.

Roles of reflex activity and co-contraction during assessments of spasticity of the knee flexor and knee extensor muscles in children with cerebral palsy and different functional levels. Pierce SR, Barbe MF, Barr AE, et al. 2008;88:1124-1134.

General

Pediatric physical therapists' perceptions of their training in assistive technology. Long TM, Perry DF. 2008;88:629-639.

Treatment

Locomotor training using body-weight support on a treadmill in conjunction with ongoing physical therapy in a child with severe cerebellar ataxia. Cernak K, Stevens V, Price R, Shumway-Cook A. 2008;88:88-97.

Time series analysis of spontaneous upper-extremity movements of premature infants with brain injuries. Ohgi S, Morita S, Loo KK, Mizuike C. 2008;88:1022-1033.

- Invited commentary. Fetters L, Scholz JP. 2008;88:1034-1036.
- Author response. Ohgi S, Morita S, Loo KK, Mizuike C. 2008;88:1037-1038.

Peer-Reviewed Publications

Editorial: New author service: PTJ makes PubMed Central deposits. Editorial Office. 2008;88:699-700.

Editorial: Still "not satisfied"... yet. Craik RL, Maher C. 2008;88:423-425.

- Correction. 2008;88:688.

Editorial: The 2008/2009 editorial board. Craik RL. 2008;88:986-988.

Journal publication productivity in academic physical therapy programs in the United States and Puerto Rico from 1998 to 2002. Richter RR, Schlomer SL, Krieger MM, Siler WL. 2008;88:376-386.

- Letter to the editor. Warden SJ. 2008;88:538-539.
- Letter to the editor. Maher C. 2008;88:539.
- Letter to the editor. Bennett SE, Ohtake PJ. 2008;88:539-540.
- Author response. Richter RR, Schlomer SL, Krieger MM, Siler W. 2008;88:540-541.
- Letter to the editor. Harris SR. 2008;88:791.

Pharmacology

Exercise and glycemic control in diabetes: benefits, challenges, and adjustments to pharmacotherapy. Gulve EA. 2008;88:1297-1321.

Phototherapy

General

Intertester reliability and validity of motion assessments during lumbar spine accessory motion testing. Landel R, Kulig K, Fredericson M, et al. 2008;88:43-49.

- Letter to the editor. Abbott JH. 2008;88:536.
- Author response. Landel R, Kulig K, Fredericson M, et al. 2008;88:536-537.

Physical Fitness

Clinical measures of physical fitness predict insulin resistance in people at risk for diabetes. Chen CN, Chuang LM, Wu YT. 2008;88:1355-1364.

Physical Therapy Profession

International

Direct access to physical therapy in the Netherlands: results from the first year in community-based physical therapy. Leemrijse CJ, Swinkels ICS, Veenhof C. 2008;88:936-946.

Patient Services

Economic realities associated with diabetes care: opportunities to expand delivery of physical therapist services to a vulnerable population. Cohn R. 2008;88: 1417-1424.

Professional Issues

Editorial: Climbing out of our silos to improve practice. Craik RL. 2008;88:555-558.

Editorial: Does clinical education need a *series* of tools to assess success? Craik RL. 2008;88: 1106-1108.

Editorial: People with diabetes: a population desperate for movement. Mueller MJ. 2008; 88:1250-1253.

Meaning of caring to 7 novice physical therapists during their first year of clinical practice. Greenfield BH, Anderson A, Cox B, Tanner MC. 2008;88:1154-1166.

Physical activity and diabetes: opportunities for prevention through policy. Deshpande AD, Dodson EA, Gorman I, Brownson RC. 2008;88:1425-1435.

Posture

General

Impact of health perception, balance perception, fall history, balance performance, and gait speed on walking activity in older adults. Talkowski JB, Brach JS, Studenski S, Newman AB. 2008;88:1474-1481.

Increased risk of falling in older community-dwelling women with mild cognitive impairment. Liu-Ambrose TY, Ashe MC, Graf P, et al. 2008;88:1482-1491.

Tests and Measurements

Use of the Berg Balance Scale for predicting multiple falls in community-dwelling elderly people: a prospective study. Muir SW, Berg K, Chesworth B, Speechley M. 2008;88:449-459.

- Invited commentary. Studenski SA. 2008;88:460-461.
- Correction. 2008;88:688.

Professional-Patient Relations

Health care providers' knowledge, attitudes, and self-efficacy for working with patients with spinal cord injury who have diverse sexual orientations. Burch A. 2008;88: 191-198.

Pulmonary

Chronic Obstructive Pulmonary Disease

Estimation of thigh muscle mass with magnetic resonance imaging in older adults and people with chronic obstructive pulmonary disease. Mathur S, Takai KPR, MacIntyre DL, Reid D. 2008;88: 219-230.

General

Breathing: a sign of life and a unique area for reflection and action. Ekerholt K, Bergland A. 2008;88:832-840.

- Letter to the editor. Taylor MJ. 2008;88:972.

Tests and Measurements

Cardiorespiratory capacity after weight-supported treadmill training in patients with traumatic brain injury. Mossberg KA, Orlander EE, Norcross JL. 2008;88:77-87.

R

Rehabilitation

After partial knee replacement, patients can kneel, but they need to be taught to do so: a single-blind randomized controlled trial. Jenkins C, Barker KL, Pandit H, et al. 2008;88:1012-1021.

Assessing recovery and establishing prognosis following total knee arthroplasty. Kennedy DM, Stratford PW, Riddle DL, et al. 2008;88:22-32.

Balance and eye movement training to improve gait in people with progressive supranuclear palsy: a quasi-randomized clinical trial. Zampieri C, Di Fabio RP. 2008;88: 1460-1473.

Changes in functional walking distance and health-related quality of life after gastric bypass surgery. Tompkins J, Bosch PR, Chenowith R, et al. 2008;88:928-935.

Computerized visual feedback: an adjunct to robotic-assisted gait training. Banz R, Bolliger M, Colombo G, et al. 2008;88: 1135-1145.

Dose estimation and surveillance of mechanical loading interventions for bone loss after spinal cord injury. Dudley-Javoroski S, Shields RK. 2008;88:387-396.

Effectiveness of an inpatient multidisciplinary rehabilitation program for people with Parkinson disease. Ellis T, Katz DI, White DK, et al. 2008;88:812-819.

Functional outcomes and quality of life after tumor-related hemipelvectomy. Beck LA, Einertson MJ, Winemiller MH, et al. 2008;88:916-927.

Ilial anterior rotation hypermobility in a female collegiate tennis player. Vaughn HT, Nitsch W. 2008;88: 1578-1590.

Inflammatory myelopathies and traumatic spinal cord lesions: comparison of functional and neurological outcomes. Scivoletto G, Cosentino E, Mammone A, Molinari M. 2008;88:471-484.

Influence of feedback schedule in motor performance and learning of a lumbar multifidus muscle task using rehabilitative ultrasound imaging: a randomized clinical trial. Herbert WJ, Heiss DG, Basso DM. 2008;88:261-269.

Physical therapy intervention for people with Huntington disease. Busse ME, Khalil H, Quinn L, Rosser AE. 2008;88:820-831.

Predictors of physical therapy clinic performance in the treatment of patients with low back pain syndromes. Resnik L, Liu D, Mor V, Hart DL. 2008;88:989-1004.

- Invited commentary. Freburger JK, Mielenz TJ. 2008;88:1004-1006.
- Invited commentary. Fritz JM. 2008;88:1007-1009.

- Author response. Resnik L, Hart DL. 2008;88:1009–1011.

Time series analysis of spontaneous upper-extremity movements of premature infants with brain injuries. Ohgi S, Morita S, Loo KK, Mizuike C. 2008;88:1022–1033.

- Invited commentary. Fetters L, Scholz JP. 2008;88:1034–1036.
- Author response. Ohgi S, Morita S, Loo KK, Mizuike C. 2008;88:1037–1038.

Usefulness of the Berg Balance Scale in stroke rehabilitation: a systematic review. Blum L, Korner-Bitensky N. 2008;88:559–566.

Use of a low-cost, commercially available gaming console (Wii) for rehabilitation of an adolescent with cerebral palsy. Deutsch JE, Borbely M, Filler J, et al. 2008;88:1196–1207.

Use of protection motivation theory, affect, and barriers to understand and predict adherence to outpatient rehabilitation. Grindley EJ, Zizzi SJ, Nasypany AM. 2008;88:1529–1540.

- Invited commentary. Thorstensson CA. 2008;88:1541–1543.
- Author response. Grindley EJ, Zizzi SJ, Nasypany AM. 2008;88:1543–1544.

Research

Benchmarking physical therapy clinic performance: statistical methods to enhance internal validity when using observational data. Resnik L, Liu D, Hart DL, Mor V. 2008;88:1078–1087.

Editorial: Still "not satisfied" . . . yet. Craik RL, Maher C. 2008;88:423–425.

Scales to assess the quality of randomized controlled trials: a systematic review. Armijo Olivo S, Macedo LG, Gadotti IC, et al. 2008;88:156–175.

S

Spinal Cord Injuries

Dose estimation and surveillance of mechanical loading interventions for bone loss after spinal cord injury. Dudley-Javoroski S, Shields RK. 2008;88:387–396.

Health care providers' knowledge, attitudes, and self-efficacy for working with patients with spinal cord injury who have diverse sexual orientations. Burch A. 2008;88:191–198.

Inflammatory myelopathies and traumatic spinal cord lesions: comparison of functional and neurological outcomes. Scivoletto G, Cosentino E, Mammone A, Molinari M. 2008;88:471–484.

Locomotor training restores walking in a nonambulatory child with chronic, severe, incomplete cervical spinal cord injury. Behrman AL, Nair PM, Bowden MG, et al. 2008;88:580–590.

- Invited commentary. Edgerton VR. 2008;88:590–591.
- Invited commentary. McEwen IR. 2008;88:592–594.
- Author response. Behrman AL, Nair PM, Bowden MG, et al. 2008;88:594–595.

T

Tests and Measurements

Functional

Associations between physical examination and self-reported physical function in older community-dwelling adults with knee pain. Wood L, Peat G, Thomas E, et al. 2008;88:33–42.

[A comparison of a modified Oswestry Low Back Pain Disability Questionnaire and the Quebec Back Pain Disability Scale. Fritz JM, Irrgang JJ. 2001;81:776–788.]

- Correction. 2008;88:138–139.

Interrater reliability of the Extended ICF Core Set for stroke applied by physical therapists. Starrost K, Geyh S, Trautwein A, et al. 2008;88:841–851.

- Invited commentary. Jette AM. 2008;88:851–853.
- Author response. Geyh S, Stucki G, Cieza A. 2008;88:854–856.

Keitel Functional Test for patients with rheumatoid arthritis: translation, reliability, validity, and responsiveness. Holm B, Jacobsen S, Skjodt H, et al. 2008;88:664–678.

Measuring deterioration in International Classification of Functioning domains of people with multiple sclerosis who are ambulatory. Paltamaa J, Sarasoja T, Leskinen E, et al. 2008;88:176–190.

Modified total-body recumbent stepper exercise test for assessing peak oxygen consumption in people with chronic stroke. Billinger SA, Tseng BY, Kluding PM. 2008;88:1188–1195.

Physical functioning before and after total hip arthroplasty: perception and performance. van den Akker-Scheek I, Zijlstra W, Groothoff JW, et al. 2008;88:712–719.

Scale for Contraversive Pushing: cutoff scores for diagnosing "pusher behavior" and construct validity. Baccini M, Paci M, Nannetti L, et al. 2008;88:947–955.

Usefulness of the Berg Balance Scale in stroke rehabilitation: a systematic review. Blum L, Korner-Bitensky N. 2008;88:559–566.

Use of the Berg Balance Scale for predicting multiple falls in community-dwelling elderly people: a prospective study. Muir SW, Berg K, Chesworth B, Speechley M. 2008;88:449–459.

- Invited commentary. Studenski SA. 2008;88:460–461.
- Correction. 2008;88:688.

General

Factors influencing the use of outcome measures for patients with low back pain: a survey of New Zealand physical therapists. Copeland JM, Taylor WJ, Dean SG. 2008;88:1492–1505.

Hierarchical properties of the motor function sections of the Fugl-Meyer Assessment Scale for people after stroke: a retrospective study. Crow JL, Harmeling-van der Wel BC. 2008;88:1554–1567.

Interrater reliability of the Movement Assessment Battery for Children. Smits-Engelsman BCM,

Fiers MJ, Henderson SE, Henderson L. 2008;88:286-294.

Predictive value of the Western Ontario and McMaster Universities Osteoarthritis Index for the amount of physical activity after total hip arthroplasty. Wagenmakers R, Stevens M, van den Akker-Scheek I, et al. 2008;88:211-218.

Reference curves for the Gross Motor Function Measure: percentiles for clinical description and tracking over time among children with cerebral palsy. Hanna SE, Bartlett DJ, Rivard LM, Russell DJ. 2008;88:596-607.

Temporal and spatial characteristics of gait during performance of the Dynamic Gait Index in people with and people without balance or vestibular disorders. Marchetti GF, Whitney SL, Blatt PJ, et al. 2008;88:640-651.

Test-retest reliability and minimal detectable change on balance and ambulation tests, the 36-Item Short-Form Health Survey, and the Unified Parkinson Disease Rating Scale in people with parkinsonism. Steffen T, Seney M. 2008;88:733-746.

- Letter to the editor. Stratford PW. 2008;88:888-890.
- Author response. Steffen TM, Seney M. 2008;88:891.

U

Ultrasound

Influence of feedback schedule in motor performance and learning of a lumbar multifidus muscle task using rehabilitative ultrasound imaging: a randomized clinical trial. Herbert WJ, Heiss DG, Basso DM. 2008;88:261-269.

Noncontact ultrasound therapy for adjunctive treatment of nonhealing wounds: retrospective analysis. Bell AL, Cavorsi J. 2008;88:1517-1524.

- Invited commentary. Robertson VJ. 2008;88:1524-1526.
- Author response. Bell AL, Cavorsi J. 2008;88:1526-1528.

Use of transabdominal ultrasound imaging in retraining the pelvic-floor muscles of a woman postpartum. Ariail A, Sears T, Hampton E. 2008;88:1208-1217.

Variability in effective radiating area at 1 MHz affects ultrasound treatment intensity. Straub SJ, Johns LD, Howard SM. 2008;88:50-57.

- Invited commentary. Robertson VJ. 2008;88:58-61.
- Author response. Straub SJ, Johns LD, Howard SM. 2008;88:61-62.

Upper Extremity

General

Modified constraint-induced therapy in chronic stroke: results of a single-blinded randomized controlled trial. Page SJ, Levine P, Leonard A, et al. 2008;88:333-340.

- Letter to the editor. Wolf SL. 2008;88:680-684.
- Author response. Page S, Levine P. 2008;88:684-688.

Reproducibility and minimal detectable change of three-dimensional kinematic analysis of reaching tasks in people with hemiparesis after stroke. Wagner JM, Rhodes JA, Patten C. 2008;88:652-663.

Time series analysis of spontaneous upper-extremity movements of premature infants with brain injuries. Ohgi S, Morita S, Loo KK, Mizuike C. 2008;88:1022-1033.

- Invited commentary. Fetters L, Scholz JP. 2008;88:1034-1036.
- Author response. Ohgi S, Morita S, Loo KK, Mizuike C. 2008;88:1037-1038.

Shoulder

Effects of Pilates exercises on shoulder range of motion, pain, mood, and upper-extremity function in women living with breast cancer: a pilot study. Keays KS, Harris SR, Lucyshyn JM, MacIntyre DL. 2008;88:494-510.

V

Vestibular System

Temporal and spatial characteristics of gait during performance of the Dynamic Gait Index in people with and people without balance or vestibular disorders. Marchetti GF, Whitney SL, Blatt PJ, et al. 2008;88:640-651.